

*Citation for published version:*

Mounce, RCP, Sansom, R & Wills, MA 2016, 'Sampling diverse characters improves phylogenies: craniodental and postcranial characters of vertebrates often imply different trees', *Evolution*, vol. 70, no. 3, pp. 666-686.  
<https://doi.org/10.1111/evo.12884>

*DOI:*

[10.1111/evo.12884](https://doi.org/10.1111/evo.12884)

*Publication date:*

2016

*Document Version*

Peer reviewed version

[Link to publication](#)

## University of Bath

### Alternative formats

If you require this document in an alternative format, please contact:  
[openaccess@bath.ac.uk](mailto:openaccess@bath.ac.uk)

#### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

#### Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

**Sampling diverse characters improves phylogenies:  
Craniodental and postcranial characters of vertebrates often  
imply different trees**

**Ross C. P. Mounce<sup>1</sup>, Robert Sansom<sup>2</sup> and Matthew A. Wills<sup>1\*</sup>**

*<sup>1</sup>The Milner Centre for Evolution, Department of Biology and Biochemistry, The University of  
Bath, The Avenue, Claverton Down, Bath  
BA2 7AY, UK; E-mail: m.a.wills@bath.ac.uk*

*<sup>2</sup>Faculty of Life Sciences, University of Manchester, Michael Smith Building, Oxford Road,  
Manchester, M13 9PT*

Words in main text : 6588 (including abstract)

Figures : 5

Tables : 2

**\*Correspondence:** Matthew A. Wills  
Department of Biology and Biochemistry  
The University of Bath  
The Avenue  
Claverton Down  
Bath BA2 7AY  
United Kingdom  
  
Telephone: 01225 383504  
Fax: 01225 386779  
E-mail: m.a.wills@bath.ac.uk

# Sampling diverse characters improves phylogenies: Craniodental and postcranial characters of vertebrates often imply different trees

Ross C. P. Mounce<sup>1</sup>, Robert Sansom<sup>2</sup> and Matthew A. Wills<sup>1</sup>

<sup>1</sup>*The Milner Centre for Evolution, Department of Biology and Biochemistry, The University of Bath,  
The Avenue, Claverton Down, Bath BA2 7AY, UK; E-mail: m.a.wills@bath.ac.uk*

<sup>2</sup>*Faculty of Life Sciences, University of Manchester, Michael Smith Building, Oxford Road,  
Manchester, M13 9PT*

Morphological cladograms of vertebrates are often inferred from greater numbers of characters describing the skull and teeth than from postcranial characters. This is either because the skull is believed to yield characters with a stronger phylogenetic signal (i.e., contain less homoplasy), because morphological variation therein is more readily atomized, or because craniodental material is more widely available (particularly in the palaeontological case). An analysis of 85 vertebrate datasets published between 2000 and 2013 confirms that craniodental characters are significantly more numerous than postcranial characters, but finds no evidence that levels of homoplasy differ in the two partitions. However, a new partition test based on tree-to-tree distances (as measured by Robinson Foulds metric) rather than tree length reveals that relationships inferred from the partitions are significantly different about one time in three, much more often than expected. Such differences may reflect divergent selective pressures in different body regions, resulting in different localized patterns of homoplasy. Most systematists attempt to sample characters broadly across body regions, but this is not always possible. We conclude that trees inferred largely from either craniodental or postcranial characters in isolation may differ significantly from those that would result from a more holistic approach. We urge the latter.

74

75 Despite the increasing importance of molecular and genomic data over the last two decades,  
76 morphology still makes an invaluable contribution to vertebrate phylogenetics, and is the only  
77 suitable source of data for palaeontological phylogenies (Asher and Müller, 2012; Hillis and Wiens,  
78 2000). Levels of morphological homoplasy in vertebrate groups are generally lower than those  
79 amongst their invertebrate counterparts (Hoyal Cuthill *et al.* 2010), suggesting that the signal  
80 quality for vertebrates is relatively high. Morphological systematists usually seek to code as many  
81 valid characters from as wide a selection of organs and body regions as possible, and to analyse  
82 these simultaneously, typically citing the principle of total evidence (Kluge, 1989). Heterogeneity in  
83 the performance of characters is often quantified in retrospect, and can also be utilised formally to  
84 increase stability in various *post hoc* weighting schemes (Farris 1969; Goloboff 2014; Goloboff *et*  
85 *al.* 2008a). However, coded matrices are usually analysed holistically rather than in partitions  
86 defined *a priori*. This is not least because hidden support (the presence of weak signals within  
87 putative partitions that become reciprocally reinforcing, and therefore the dominant signal when all  
88 characters are considered) is most readily identified in this way (Gatesy *et al.* 1999; Gatesy and  
89 Arctander 2000). It is therefore conceded that different body regions can be subject to different  
90 levels and patterns of homoplasy (Gaubert *et al.* 2005), such that anatomical subsets of characters  
91 can yield significantly different trees. This is especially true when there is strong functional  
92 selection in particular organ systems, leading to convergence (Ji *et al.* 1999; Kivell *et al.* 2013;  
93 Tseng *et al.*, 2011).

94 Few cladistic studies have analysed the performance of morphological characters within  
95 partitions explicitly (e.g., Sánchez-Villagra and Williams, 1998; Rae, 1999; Gould, 2001; Song and  
96 Bucheli, 2010). While analyses may concentrate upon characters of particular types or from  
97 particular organ systems, it is still rare that trees are inferred explicitly from subsets of these data  
98 (but see Rae, 1999; Vermeij and Carlson, 2000; O'Leary *et al.*, 2003; Poyato-Ariza, 2003; Diogo,  
99 2004; Young, 2005; Clarke and Middleton 2008; Smith, 2010; Farke *et al.*, 2011). Notable  
100 exceptions are taphonomic studies that investigate the effects of omitting volatile, soft-part



101 characters with a low fossilisation potential (Sansom *et al.* 2010, 2011; Sansom and Wills, 2013;  
102 Pattinson *et al.*, 2014; Sansom 2015).

103       Where levels of homoplasy across morphological character partitions are considered at all,  
104 these are usually investigated by comparing distributions of the consistency index (ci: Kluge and  
105 Farris, 1969) (Fig. 1). The ci is given simply as the minimum possible number of state changes  
106 (states minus characters) divided by the most parsimonious tree length. More appropriate functions  
107 underpin most *post hoc* weighting schemes (Goloboff 2014; Goloboff *et al.* 2008a). In the case of  
108 insects, Song and Bucheli (2010) found male genital characters to be less homoplastic than  
109 characters coding other aspects of form. In brachiopods, Leighton and Maples (2002) revealed that  
110 shell characters are significantly more homoplastic than those describing internal anatomy (a  
111 disconcerting finding in a group whose fossil record consists largely of hard parts, and in which  
112 phylogeny is inferred almost exclusively with recourse to such characters). Similarly, in hedgehogs,  
113 Gould (2001) reported significantly higher consistency indices for dental characters compared with  
114 those describing other aspects of anatomy. However, she also noted that the optimal trees inferred  
115 from the dental characters alone were seriously at odds with those inferred from the other  
116 characters. Kangas *et al.* (2004) noted that many morphological characters of mammalian teeth  
117 have exceptionally strong developmental interdependence, all being under the correlated control of  
118 a small number of genes. Most ambitiously, Sanchez-Villagra and Williams (1998) compared the  
119 consistency indices between dental, cranial and postcranial character partitions of eight  
120 mammalian datasets, but reported no significant differences. We note that the relative performance  
121 of particular characters and character partitions may be contingent, especially with respect to the  
122 taxonomic hierarchical level of study. Hence, characters entailing little or no homoplasy for shallow  
123 branches may exhibit greater homoplasy when considered at deeper levels. Goloboff *et al.* (2010)  
124 have proposed analytical approaches that take account of this variability in the evolutionary lability  
125 of characters along branches.

126       The dominant practice of total evidence analysis for morphology contrasts with the more  
127 qualified approach that was sometimes adopted with molecular data (Bull *et al.*, 1993; Felsenstein,  
128 1988; Pamilo and Nei, 1988; Maddison, 1997; Nichols, 2001; Degnan and Rosenberg, 2009).

129 Molecular systematists have debated the relative merits of partitioned versus combined analyses,  
130 although the consensus has emerged in favour of the latter (Gatesy *et al.*, 1999; Gatesy and  
131 Baker, 2005; Kjer and Honeycutt, 2007; Thompson *et al.*, 2012). Historically, the issue of  
132 morphological versus molecular incongruence has been more to the fore (Gatesy and Arctander,  
133 2000), motivated by striking examples of conflict between molecular and morphological  
134 cladograms in some groups (Mickevich and Farris, 1981; Bledsoe and Raikow, 1990; Bremer,  
135 1996; Poe, 1996; Baker *et al.*, 1998; Hillis and Wiens, 2000; Wiens and Hollingsworth, 2000;  
136 Jenner, 2004; Draper *et al.*, 2007; Springer *et al.*, 2007; Pisani *et al.*, 2007; Near 2003; Mayr,  
137 2011), although see Lee and Camens (2009).

138

### 139 **WHY EXAMINE THE CONGRUENCE OF CRANIODENTAL AND POSTCRANIAL PARTITIONS?** 140

141 Some systematists hold that craniodental and postcranial characters convey signals of differing  
142 quality (Ward, 1997; Collard *et al.*, 2001; Naylor and Adams, 2001; Finarelli and Clyde, 2004).  
143 However, the evidence for this is piecemeal and largely anecdotal. Many practitioners take a more  
144 holistic approach to sampling characters (Sánchez-Villagra and Williams, 1998), sampling densely  
145 from as many anatomical regions as possible. However, even where potential characters are  
146 reasonably homogenously distributed throughout the body, “certain body regions and organs still  
147 hold a considerable mystique for taxonomists as classificatory tools, while others are neglected”  
148 (Sokal and Sneath, 1963; page 85). Arratia (2009) noted that actinopterygian systematists focus on  
149 cranial characters, despite rich seams of underexploited data within the fin rays and fulcra. Murray  
150 and Vickers-Rich (2004) suggested that the crania and mandibles of birds often provide the most  
151 informative characters because of their structural complexity. Similarly, Cardini and Elton (2008)  
152 demonstrated that characters of the chondrocranium were most informative in studies of  
153 *Cercopithecus* monkeys, and suggested that this might apply across primates and perhaps across  
154 all mammals. Lastly, Ruta and Bolt (2008) found that characters of the lower jaws of  
155 temnospondyls recovered many of the same relationships as those inferred from a more holistic  
156 data set.

157           Studies of extinct organisms *necessarily* focus on those characters capable of fossilisation  
158 (typically shells and bones). In vertebrates, disproportionate numbers of characters are often  
159 coded from the most recalcitrant skeletal elements, notably teeth in mammals (Billet, 2011;  
160 Alejandra Abello, 2013). More generally, there is a tendency for soft part characters to resolve as  
161 more derived apomorphies than those characters with a higher preservation potential. Removal of  
162 soft characters therefore results in preferential 'stemward slippage' across animal groups (Sansom  
163 *et al.*, 2010, 2011; Sansom and Wills, 2013); the phenomenon whereby taxa resolve closer to the  
164 root of the tree than they otherwise would.

165           In this study, we apply a variety of methods to explore differences in the strength and  
166 nature of phylogenetic signals in craniodental and postcranial partitions of 85 published vertebrate  
167 datasets. We address the following questions: 1. Do levels of homoplasy in craniodental character  
168 partitions differ from those in postcranial character partitions (Sanchez-Villagra and Williams,  
169 1998), and are any observed differences more than simply a function of differing numbers of  
170 characters within these partitions? We quantify this using conventional indices of homoplasy  
171 (Kluge and Farris, 1969; Archie, 1996) modelled with respect to data set parameters. 2. Is there  
172 more conflict between craniodental and postcranial characters than we would expect for random  
173 partitions, and do craniodental and postcranial characters support significantly different trees as a  
174 result? We investigate this using the incongruence length difference test (Mickeyevich and Farris,  
175 1981; Farris *et al.*, 1995a,b) and a new partition homogeneity test based upon tree distance  
176 metrics rather than differences in tree lengths.

177

178

## 179 *Materials and Methods*

180

### 181 **THE DATA SETS**

182

183 Phylogenetic datasets published between 2000 and 2013 were sourced from the literature. We  
184 restricted our focus to discrete character morphological matrices composed entirely of vertebrate  
185 taxa, and analysed using equal weights maximum parsimony. Although morphological data can be

186 analysed with model-based likelihood (Lewis, 2000; Lee and Worthy, 2012) and Bayesian  
187 (Nylander *et al*, 2003; Pollitt *et al.*, 2005; Clarke and Middleton, 2008; Tsuj and Mueller, 2009;  
188 Bouchard-Cote *et al*, 2012) methods, the considerable majority of published morphological trees  
189 are generated utilising maximum parsimony. Matrices were also garnered from Brian O'Meara's  
190 *TreeBASE* mirror (O'Meara, 2009), Graeme Lloyd's collection of dinosaur matrices (Lloyd, 2009)  
191 and *MorphoBank* (O'Leary and Kaufman, 2011). We excluded matrices with fewer than eight taxa  
192 or partitions with fewer than eight parsimony-informative characters (for reasons of statistical  
193 power). We interpret craniodental characters here as those pertaining to the skull (cranium plus  
194 mandible and dentition).

195 Our resulting sample comprised 85 matrices, spanning all major vertebrate groups. A small  
196 number of these datasets contained characters that were not strictly morphological (e.g., character  
197 618 relating to habitat choice in the matrix of Spaulding *et al.*, 2009). These were removed prior to  
198 any further analysis. We also removed phylogenetically uninformative taxa within partitions using  
199 the principles of safe taxonomic reduction (Wilkinson, 1995). We additionally removed taxa with  
200 large amounts of missing data that were demonstrated empirically to obfuscate resolution within  
201 partitions, usually because they inflated search times and numbers of optimal trees to impractical  
202 levels. The number and percentage of craniodental and postcranial characters in each matrix were  
203 recorded, as well as the fraction of missing entries within these partitions. Wilcoxon signed-rank  
204 tests were used to assess the significance of differences between these medians in different  
205 groups.

206 Simply recording percentages of missing cells has limitations, as these can be distributed  
207 randomly (usually less problematic analytically) or can be concentrated within particular taxa. Such  
208 concentrations are often observed in real data sets, and particularly in matrices including fossil  
209 taxa (Cobbett *et al.*, 2007). Our pruning of taxa removed the worst of these effects. Simulations  
210 have demonstrated that it is the signal within the characters that *are* coded that is critical in  
211 determining the placement of particular taxa (Wiens, 2003ab) rather than numbers of missing cells  
212 *per se*.

213 All phylogenetic analyses were performed using *TNT* (Goloboff *et al.*, 2008b), using equally  
214 weighted parsimony. We also reproduced any assumptions regarding character order, polarity and  
215 rooting. Empirically, we determined that comprehensive searches involving 200 parsimony ratchet  
216 iterations (Nixon, 1999) and 100 drift iterations per replication, with 10 rounds of tree fusion  
217 (Goloboff, 1999) were effective at recovering the set of MPTs reported by the original authors in  
218 each case. These settings were used throughout.

219  
220 **TESTING WHETHER CRANIODENTAL AND POSTCRANIAL CHARACTERS EXHIBIT**  
221 **DIFFERENT LEVELS OF HOMOPLASY**  
222

223 *Consistency and retention indices.* – There are two intuitive ways to calculate differences in  
224 mean/median consistency indices (*ci*; Kluge and Farris, 1969) and retention indices (*ri*; Farris,  
225 1989) for characters in partitions of a dataset (*ci* and *ri* in lower case pertain to individual  
226 characters). The usual approach is to find the optimal tree or trees for all characters analysed  
227 simultaneously (the global MPT(s)) and to take mean values for characters reconstructed on  
228 this/these (Sánchez-Villagra and Williams, 1998; Song and Bucheli, 2010). However, there are  
229 theoretical partition size effects, even in the absence of differences in the levels of character  
230 conflict *within* partitions. All other things being equal, the characters within the larger partition are  
231 likely to have higher *ci* values on average (Fig. 1). The other approach is to report metrics for  
232 characters within each partition analysed independently. However, the ensemble CI and ensemble  
233 RI (and therefore *ci* and *ri* for individual characters) are influenced by data set dimensions (Archie,  
234 1989; Sanderson and Donoghue, 1989; Faith and Cranston, 1991; Klassen *et al.*, 1991): there is a  
235 strong, negative correlation between CI and the number of taxa and a weaker, negative  
236 relationship between CI and the number of characters (Archie, 1989, 1996; Archie and Felsenstein,  
237 1993).

238 There are two ways in which differences between indices can be tested. For individual data  
239 matrices, Mann-Whitney or t-tests can be applied to character *ci* and *ri* values, with the null that  
240 these have the same median or mean in the two partitions. For the more general comparison  
241 across all 85 matrices simultaneously, Wilcoxon signed ranks or paired t-tests can be used to test

242 the nulls that (either) the median/mean ci or ri in craniodental and postcranial partitions were  
243 similar, or that the median/mean CI and RI indices for the two partitions were similar.

244

245 *Homoplasy excess ratio (HER)*. – The homoplasy excess ratio (HER; Archie and Felsenstein,  
246 1993) was proposed as an adjunct to the ensemble consistency index (CI), and argued to be  
247 relatively immune to its worst shortcomings. HER is given by  $(\text{MEANNS} - L) / (\text{MEANNS} - \text{MINL})$ ,  
248 where MEANNS is the mean length of the most parsimonious trees resulting from a large sample  
249 of matrices (here 999) in which the state assignments within each character have been  
250 randomised. L is then the optimal length of the original dataset, and MINL is the minimum possible  
251 length of the dataset (number of states minus number of characters). HER was calculated for  
252 craniodental and postcranial partitions in isolation, and we then tested for differences in partitions  
253 across all 85 datasets using the Wilcoxon signed ranks test.

254

## 255 **TESTING INCONGRUENCE BETWEEN CRANIODENTAL AND POSTCRANIAL CHARACTER** 256 **PARTITIONS**

257  
258 *Incongruence length difference (ILD) test*. – To assess the significance of congruence between  
259 whole character partitions as measured by optimal tree length, the ILD test (Mickevich and Farris,  
260 1981; Farris *et al.*, 1995ab; Barker and Lutzoni, 2002) was applied to the matrices in *TNT* using  
261 999 replicates (Allard *et al.*, 1999a,b). The ILD score is given by  $L_{AB} - (L_A + L_B) / L_{AB}$ , where  $L_{AB}$  is  
262 the optimal tree length (in steps) of the simultaneous analysis of both partitions together (the total  
263 evidence analysis), and  $L_A$  and  $L_B$  are the optimal tree lengths for partitions A and B analysed  
264 independently. To determine the significance of the observed ILD score, random partitions of the  
265 same size (number of characters) as the specified partitions are also generated to yield a  
266 distribution of randomized ILD scores. Given the nature of phylogenetic data, the suitability of this  
267 test has been questioned on a variety of grounds (Dolphin *et al.* 2000; Hipp *et al.* 2004; Ramirez  
268 2006; Planet 2005, 2006). Despite this, the ILD test remains commonly used to compare the  
269 congruence of data partitions. We did not apply the arcsine transformation of Quicke *et al.* (2007)  
270 because they justified their correction on the basis of empirical and simulated *molecular* data  
271 (morphological data have different statistical properties).

272

273 **TESTING WHETHER PARTITIONS SUPPORT DIFFERENT TREES**

274

275 *The incongruence relationship difference (IRD) test: a new test of the congruence of relationships.*

276 – Much like the ILD test, this is a randomisation-based test. However, partitions are compared via  
277 the distances between the optimal trees that result from them, rather than via tree length (ILD) or a  
278 matrix-representation of topology (TILD; Wheeler, 1999). There are many possible tree-to-tree  
279 distance measures including symmetric difference (RF; Bourque, 1978; Robinson and Foulds,  
280 1981; Pattengale *et al.* 2007) quartets distance (QD; Estabrook *et al.*, 1985), nearest neighbour  
281 interchange distance (NNID; Waterman and Smith, 1978), nodal distance (Bluis and Shin, 2003),  
282 maximum agreement subtree distance (Goddard *et al.*, 1994; de Vienne *et al.*, 2007), transposition  
283 distance (Rossello and Valiente, 2006), subtree prune and regraft distance (SPR; Goloboff, 2008),  
284 and path-length difference (PLD; Zaretskii, 1965; Williams and Clifford, 1971). For reasons of  
285 familiarity (it is among the most well characterised; e.g. Steel and Penny, 1993) and ease of  
286 computation, we chose to use the symmetric difference or Robinson and Foulds distance (RF)  
287 (Fig. 2) as our measure of tree-to-tree distance. We note that all other implementations are  
288 possible. We illustrate the approach for two examples: firstly the theropod data of Ezcurra and  
289 Cuny (2007) (Fig. 3a) and secondly the mammalian data of Beck (2008) (Fig. 3b). The results from  
290 each partition are illustrated as 50% majority rule consensus trees for ease of visualization. The left  
291 hand tree in both cases is that derived from the analysis of craniodental characters alone, while the  
292 right hand tree is inferred from just the postcranial characters. The open circles indicate nodes  
293 present in one partition tree that are absent from the other, and the total number of such nodes  
294 gives the measure of RF between the trees. This value corresponds to the incongruence  
295 relationship difference ( $IRD_{MR}$  in the case of the majority rule trees illustrated).

296 All MPTs from the analysis of each partition were saved and then compared to each other  
297 in two different ways. (1) ‘Nearest neighbours’ ( $IRD_{NND}$ ) for up to 10,000 trees in each partition: the  
298 mean of the minimum distance between each tree in one set, compared with the trees in the other  
299 (and *vice versa*) (Cobbett *et al.*, 2007). (2) The distance between the 50% majority-rule consensus  
300 trees (from up to 10,000 fundamentals) for each partition ( $IRD_{MR}$ ) (Figs. 3 and 4). We then

301 generated random partitions of the original data in the original proportions, and repeated the above  
302 exercises in order to yield a distribution of randomized partition tree-to-tree distances. Distances  
303 for the original partitions were deemed significantly different from this distribution when they lay in  
304 its 5% tail. Our p-values were derived from 999 replicates. All but three (98%) of our 170 partitions  
305 yielded less than 10,000 trees; the imposition of a 10,000 bound for the remainder was a  
306 necessary limitation to restrict prohibitively long searches in poorly resolved partitions.

307

308 *Tests not implemented.* – The topological incongruence length difference (TILD) test (Wheeler,  
309 1999) is analogous to the ILD test, but is applied to a matrix representation (GIC; Farris, 1973; also  
310 known as MRP coding, Baum, 1992; Ragan, 1992) of the branching structure of a consensus of  
311 the optimal trees from the data partitions. The test appears to have limited discriminatory power  
312 and high type I error rate (Wills *et al.* 2009).

313 Rodrigo *et al.* (1993) proposed three interrelated tests to investigate differences in  
314 relationships directly. The first of these determines whether the symmetrical difference distance  
315 (RF: Robinson and Foulds, 1981) between sets of MPTs from independent analyses of the two  
316 data set partitions is distinguishable from the distribution of RF distances between a large sample  
317 of pairs of random trees. Only weak congruence between partitions is needed to pass this test. The  
318 second test of Rodrigo *et al.* (1993) compares the partitions directly, and determines if there is any  
319 overlap between the MPTs derived from the two partitions upon bootstrap resampling. This is  
320 problematic because the probability of encountering common trees changes with the bootstrap  
321 parameters (Lutzoni, 1997), especially the number of replicates (Page, 1996). The third test  
322 compares RF distances between partitions with those between trees bootstrapped from within  
323 partitions. Although a useful test, it may have limitations, particularly where the partitions of the  
324 data set are of very different sizes, and especially where the number of characters in the smaller  
325 partition is also small relative to the number of terminals. In such cases, bootstraps of the smaller  
326 partition may consistently yield poor resolution and low RF distances between trees *within* this  
327 partition (Page, 1996). The IRD<sub>NND</sub> test proposed above controls this partition size difference.

328



## 329 *Results*

330

### 331 **HOMOPLASY IN CRANIODENTAL AND POSTCRANIAL DATA PARTITIONS**

332

333 Across our sample of 85 datasets, craniodental partitions had more characters (median = 58) than  
334 postcranial partitions (median = 50) (Wilcoxon signed ranks;  $V = 2288.5$ ,  $p = 0.044$ ) (Table 1). The  
335 percentage of missing data cells was comparable in craniodental (median = 12.8%) and  
336 postcranial (median = 16.9%) partitions, although this difference was significant (paired Wilcoxon;  
337  $V = 868.5$ ,  $p = 0.006$ ). The mean ensemble consistency indices (CI) for craniodental and  
338 postcranial characters across all 85 data sets were not significantly different (paired  $t = 1.184$ ,  $p =$   
339  $0.240$ ), with postcranial partitions ( $\bar{X} = 0.564$ ) having slightly higher values than craniodental  
340 partitions ( $\bar{X} = 0.550$ ). Using the mean partition (per character) ci index across all matrices  
341 (characters optimised onto the globally optimal tree(s) for the entire matrix) revealed a non-  
342 significant difference ( $\bar{X} = 0.632$  and  $0.627$  for craniodental and postcranial partitions respectively;  
343 paired  $t = 0.450$ ,  $p = 0.654$ ). Mann-Whitney tests of craniodental and postcranial ci values *within*  
344 the 85 data sets yielded 40 significant ( $p < 0.05$ ) results (four or five might be expected). Twenty-  
345 one of these 40 had higher means (less homoplasy) for cranial characters, despite their larger  
346 partition size. A simple linear model was used to express partition CI in terms of the log of the  
347 number of taxa, the log of the number of characters and the log of the percentage of missing data  
348 (+1) across all 170 partitions. The term for missing data was not significant, but both the log of the  
349 number of taxa ( $p < 0.001$ ) and the log of the number of characters ( $p < 0.014$ ) were highly so  
350 (multiple  $R^2 = 0.458$ ,  $p = < 0.001$ ). A subsequent paired t-test of the residual CI values from this  
351 model revealed no significant difference ( $t = 0.917$ ,  $p = 0.362$ ) between craniodental and  
352 postcranial partitions. We note that other variables have been demonstrated empirically to  
353 influence CI (Donoghue and Ree, 2000; Hoyal Cuthill *et al.*, 2010).

354 Homoplasy excess ratio (HER) values (Archie 1989, 1996; Archie and Felsenstein, 1993)  
355 were similar in the craniodental ( $\bar{X} = 0.582$ ) and postcranial skeleton ( $\bar{X} = 0.571$ ) (paired  $t =$   
356  $0.621$ ,  $p = 0.537$ ). A linear model of HER in terms of the logs of numbers of characters and taxa  
357 and the percentage of missing data (+1) revealed no significant independent variables. Finally,

retention indices were significantly higher for postcranial than craniodental partitions when measured across all characters in a partition (RI; paired  $t = 2.654$ ,  $p = 0.009$ ) but not as the average of per character values within a partition (ri; paired  $t = 1.538$ ,  $p = 0.128$ ). Linear modelling of the partition RI in terms of the logs of numbers of characters and taxa and the percentage of missing data (+1) revealed no significant independent variables.

363

## 364 CONGRUENCE BETWEEN CRANIODENTAL AND POSTCRANIAL SIGNALS (ILD TESTS)

365

366 When originally described, the ILD test was used with a standard significance level of 5% (0.05). At  
367 this level, 31 of our 85 dataset partitions had significant character incongruence (Table 1). Some  
368 have advocated more stringent levels (e.g., Cunningham, 1997): with  $p < 0.010$ , we still rejected  
369 the null for 23 of our datasets. We note that the correlation between ILD  $p$ -values and the  
370 percentage of missing data within a data set was not significant ( $r = -0.115$ ,  $p = 0.125$ ), although  
371 there was a significant correlation between ILD  $p$ -values and the *difference* in the percentage of  
372 missing data cells in the two partitions ( $r = -0.161$ ,  $p = 0.031$ ). When culling our matrices to those  
373 with a difference of just 5% or less in the fraction of missing cells in the two partitions ( $n = 40$ ), we  
374 still observed 11 data sets with an ILD significant at  $p < 0.050$  (a similar rate of rejection:  $G =$   
375  $2.653$ ,  $p = 0.103$ ).

376 Logistic regression was used to model the binary outcome of the ILD test (significant or not  
377 at  $p < 0.05$ ) as a function of log of the number of taxa, log of the number of characters, the  
378 imbalance in number of characters between partitions (as a percentage of the total number), the  
379 percentage of missing entries in the data set, and the log of the imbalance in the percentage of  
380 missing entries between partitions. Terms for the number of characters ( $p < 0.001$ ) and the  
381 imbalance in character numbers ( $p = 0.021$ ) were retained in the minimum adequate model  
382 selected by the progressive deletion of non-significant terms ( $p > 0.05$ ).

383

## 384 THE SIMILARITY OF RELATIONSHIPS IMPLIED BY PARTITIONS

385

386 Across all 85 data sets, 27 had significantly incongruent relationships (IRD<sub>NND</sub>) at  $p < 0.05$ , of  
387 which 14 were also significant at  $p < 0.01$  (Table 1). Correlation between  $p$ -values and the

388 difference in the percentage of missing data between partitions was not significant ( $\tau = -0.012$ ,  $p =$   
389  $0.876$ ). Moreover, the rate of rejection of the null at  $p < 0.05$  was similar for the culled ( $n = 40$ ) data  
390 set and those matrices ( $n = 45$ ) with more than 5% difference in missing data between partitions  
391 (eleven and sixteen significant results respectively) ( $G = 0.637$ ,  $p = 0.425$ ). Logistic regression of  
392 the binary outcome of the  $IRD_{NND}$  test ( $p < 0.05$  or otherwise) as above yielded no significant terms  
393 in the minimum adequate model.

394 The correlation between  $p$ -values for the nearest neighbour and majority rule variants of the  
395  $IRD$  test was highly significant ( $p < 0.001$ ) but not particularly tight ( $\tau = 0.387$ ). The latter offers a  
396 very imprecise proxy for the distances measured by nearest neighbours, and we do not advocate  
397 its use.

398

## 399 **HIGHER TAXONOMIC DIFFERENCES IN CONGRUENCE**

400

401 Partitioning datasets into six broad and inclusive taxonomic groups (Aves, other Ornithodira,  
402 Synapsida, other reptiles, amphibians (including early tetrapods), and fishes) revealed some  
403 significant differences (Fig. 5). In particular, synapsids and amphibians were less likely to have  
404 congruent partitions according to the  $ILD$  test than other groups ( $G = 11.808$ ,  $p = 0.038$ ) (Table 2).  
405 However, there were significant differences in the log of the number of characters across groups ( $F$   
406  $= 3.095$ ,  $p = 0.013$ ), largely accounted for by the contrast between synapsids and fishes (Tukey  
407  $HSD$ ,  $p = 0.003$ ). Taxonomic group was not significant as a factor in logistic regression models,  
408 and the residuals from models omitting taxonomic group membership retained no differences  
409 between groups (Kruskal-Wallis  $\chi$ -squared  $= 2.976$ ,  $p = 0.704$ ). A broadly similar taxonomic pattern  
410 was observed for the  $IRD_{NND}$  test, although differences in frequencies of null rejection across  
411 higher taxa were not significant ( $G = 4.283$ ,  $p = 0.510$ ).

412

## 413 *Discussion*

414

## 415 **CRANIODENTAL AND POSTCRANIAL PARTITIONS CONTAIN SIMILAR LEVELS OF**

## 416 **HOMOPLASY**

417  
418 It is well known that the CI is negatively correlated with the number of taxa in a matrix (Archie,  
419 1989; Sanderson and Donoghue, 1989; Faith and Cranston, 1991; Klassen *et al.*, 1991), but it also  
420 has a weaker, negative relationship with the number of characters (Archie, 1989, 1996; Archie and  
421 Felsenstein, 1993). Despite the greater number of craniodental characters than postcranial  
422 characters across our sample of data sets, we found no significant difference in CI (paired  $t =$   
423  $1.184$ ,  $p = 0.240$ ). Unsurprisingly, when modelling out both data matrix dimensions (numbers of  
424 taxa and characters) and the percentage of missing data (+1) across all 170 partitions, residual CI  
425 values were even more similar (for residuals:  $t = 0.917$ ,  $p = 0.362$ ). Homoplasy excess ratio (HER)  
426 values were also similar in the craniodental ( $\bar{x} = 0.582$ ) and postcranial skeleton ( $\bar{x} = 0.571$ )  
427 (paired  $t = 0.621$ ,  $p = 0.537$ ).

428 We note that the absence of a clear difference between craniodental and postcranial levels  
429 of homoplasy does not necessarily imply that *additional* characters of equivalent phylogenetic  
430 informativeness can be garnered from the two partitions with comparable ease. One partition may  
431 have been exhausted with considerable care, the other not. Our conclusions therefore necessarily  
432 relate to the *coded* data. We also note that a high CI within a partition could be the result of a  
433 strong phylogenetic signal, or the developmental non-independence of characters. Distinguishing  
434 between these causes requires detailed developmental and underpinning genetic knowledge,  
435 which are often unavailable.

436

## 437 **PARTITIONS HAVE INCONGRUENT SIGNALS MORE OFTEN THAN WE EXPECT**

438

439 Our ILD test results demonstrate that our partitions are incongruent about one time in three: 31  
440 from 85 datasets. Assuming a significance level (false positive rate) of 5%, we would expect four or  
441 five datasets to be significantly incongruent by chance. Significant incongruence is therefore  
442 detected across our sample of datasets (binomial test  $p < 0.001$ , assuming a 5% false positive  
443 error rate), although this is partly accounted for by differences in partition parameters. We make no  
444 inferences concerning the overall quality of individual data sets on the strength of these results,

445 and note that partitions were imposed by us in each case (rather than reflecting distinctions made  
446 by the original authors).

447

## 448 **CRANIODENTAL AND POSTCRANIAL PARTITIONS OFTEN IMPLY SIGNIFICANTLY** 449 **DIFFERENT RELATIONSHIPS**

450

451 Results from the incongruence relationship difference (IRD<sub>NND</sub>) tests were broadly similar to  
452 those from the ILD test: 32% of data sets yielded significantly different trees from the two partitions.  
453 As with the ILD test, we would expect just three or four data sets (5%) to reject the null by chance  
454 (a highly significant difference: binomial test  $p < 0.001$ ). Our empirical sample suggests that the  
455 IRD<sub>NND</sub> test using the Robinson Foulds distance is less likely to yield a significant result than the  
456 ILD, and is less susceptible to differences in partition size and differences in levels of missing data.  
457 The IRD<sub>NND</sub> therefore has certain advantages over the ILD, and also offers a more intuitive index of  
458 congruence (i.e., one based directly on differences in topological branching structure rather than  
459 differences in tree length). We note, however, that the IRD<sub>NND</sub> is insensitive to differences in branch  
460 lengths. Many authors have identified difficulties with the incongruence length difference test (ILD)  
461 (Dolphin *et al.* 2000; Hipp *et al.* 2004; Ramirez 2006; Planet 2005, 2006) and we do not repeat  
462 these here.

463 As with all similar metrics, the IRD is to some extent arbitrary (Wheeler, 1999). Different  
464 indices of tree-to-tree distances will yield different distances and p-values, and the IRD can be  
465 implemented with many such measures. For example, the Robinson Foulds distance penalises  
466 distant and shallow branch transpositions much more strongly than close and shallow  
467 transpositions, whereas the maximum agreement subtree distance (Finden and Gordon, 1985), for  
468 example, makes only a marginal distinction between these two cases. Similarly, the maximum  
469 agreement subtree distance is less sensitive to the depth of the transposition (Cobbett *et al.*,  
470 2007). Alternative metrics will have other, and perhaps more desirable, properties (Lin *et al.*, 2011).  
471 Our choice of the Robinson Foulds distance here was partly pragmatic, as it is computationally less  
472 demanding than most alternatives.

473

## 474 WHAT DOES PARTITION INCONGRUENCE IMPLY?

475

476

477

478

479

480

481

482

483

484

485

486

487

488

489

490

491

492

493

494

495

496

497

498

499

500

501

502

In studying the evolution of form, it is now relatively common to recognize anatomical modules (Mitteroecker and Bookstein, 2007, 2008; Klingenberg, 2008; Cardini and Elton, 2008; Lü *et al.* 2010; Hopkins and Lidgard, 2012; Cardini and Polly, 2013; Goswami *et al.*, 2013, 2015). These are regions of the body (or suites of landmarks) within which morphological changes are strongly correlated through evolutionary time, but between which there is significantly less coordination. Different selective forces may operate on these modules or components of the mosaic (Gould, 1977; Maynard Smith, 1993; Kemp, 2005; Lü *et al.*, 2010), and they may therefore exhibit different evolutionary rates and trends (Mitteroecker and Bookstein, 2007, 2008; Klingenberg, 2008). In the context of phylogenetic characters, differing pressures on modules may favour particular patterns of convergence and homoplasy, and therefore suites of characters that imply different trees (Clarke and Middleton, 2008). The skull of many tetrapod groups has often been regarded as biomechanically and functionally somewhat independent of the rest of the skeleton (Ji *et al.*, 1999; Koski, 2007; Mitteroecker and Bookstein, 2008) hence the difficulty of making many inferences about the one from the other. We note that fishes have the lowest overall incongruence between partitions and the most similar levels of per character consistency index (ci) (Fig. 5). This may result from greater functional and biomechanical integration between the head and trunk in fishes compared with other vertebrate groups (i.e., fishes lack a functional neck) (Klingenberg, 2008; Larouche *et al.*, 2015).

We note that while anatomical modules are usually envisaged as physically proximate suites of landmarks or characters, it is possible for characters to evolve in a coordinated manner across the body as the result of particular selective pressures (Kemp, 2007). For example, adaptations for swimming, digging or flying (Gardiner *et al.*, 2011; Abourachid and Hofling, 2012; Allen *et al.*, 2013) might entail correlated suites of change across the body in a manner that would not be apparent from studying straightforward divisions into body regions (e.g., head, body and limbs). Developmental regulatory processes may also entail counterintuitive suites of coordinated character change across the body (Kharlamova *et al.* 2007; Chase *et al.* 2011), but our objective was not to test for such suites of correlations here.

503 In the most general terms, character selection and coding clearly has an impact upon  
504 inferred phylogeny. Most straightforwardly, alternative data sets for identical sets of taxa can yield  
505 different trees (Freitas and Brown, 2004; Munoz-Duran, 2011; Penz *et al.* 2013), and this argues  
506 strongly for the synthesis of all characters. More generally, systematists rightly exercise their  
507 judgement in deciding which aspects of morphology to codify as putative homologies. Wings in  
508 birds, bats and pterosaurs are not considered homologous as wings *a priori* (although they are as  
509 *limbs*) because the weight of evidence unambiguously rules this out (and in the absence of any  
510 formal analysis). In many cases, however, the decision is less straightforward, and the *a priori*  
511 omission of characters believed to be analogous or strongly homoplastic may unintentionally  
512 overlook useful signal at some level in the tree. Finally, it is often observed empirically that the  
513 tree(s) derived from a given matrix can alter markedly with the omission, reweighting or ordering of  
514 characters (Wills, 1998), such that even modest perturbations to the data yield large changes to  
515 the resulting trees.

516 In morphological phylogenetics, it is usual to combine all available data (Kluge *et al.*, 1989).  
517 Our results therefore reinforce the importance of this approach. While the patterns inferred from  
518 particular organ systems or suites of characters may be misleading (in the same way and for the  
519 same reasons that individual characters may merely introduce homoplasy and noise), combined  
520 analysis of all available characters often allows a globally strong phylogenetic signal to emerge  
521 from conflicting local homoplasy (Gatesy *et al.* 1999). We demonstrate that in vertebrate studies of  
522 this type, an exclusive focus upon characters of either the cranium or postcranium (at the expense  
523 of those of the other partition (e.g., Fitzgerald (2010) (craniodental only) and Mayr and Mourer-  
524 Chauvire (2004) (postcranial only)) will significantly influence the resultant optimal tree(s) about  
525 30% of the time, irrespective of major group. We therefore strongly advocate garnering character  
526 data intensively from all anatomical regions whenever possible.

527 When analysing fossils, it is usually impossible to sample across the same suite of  
528 characters that would ideally be coded for extant species (Wiens, 2003a,b; Cobbett *et al.*, 2007).  
529 For example, in fossil crocodyliforms, the vast majority of characters are coded from the skull (e.g.,  
530 O'Connor *et al.*, 2010; Turner and Sertich, 2010; Cau and Fanti, 2011; Hastings *et al.*, 2011;

Puertolas *et al.*, 2011) and it is difficult to be confident that we are not merely inferring a 'craniodental' tree. The only (and indirect) way to test this would be to conduct parallel analyses upon the closest living representatives of the clade. However, the (quite possibly limited) utility of this approach depends upon the phylogenetic proximity of the extant exemplars, the presumed constancy of selective pressures on putative modules through time and across clades (a big assumption: Hunt, 2008; Frazzetta 2012), and the similarity of the available coded data. A related issue in the context of fossil vertebrates is the preferential preservation of hard part characters (bones rather than muscles or other more volatile tissues). An analogous concern, therefore, is whether skeletal and soft-part characters convey a consistent phylogenetic signal (Diogo, 2004). If they do not, then this has implications for the manner in which fossil vertebrates are interpreted and analyzed (Sansom *et al.*, 2010; Sansom and Wills 2013; Pattinson *et al.*, 2014) and is an area particularly needing detailed future work.

543

## 544 *Conclusions*

545

- 546 1. Systematists typically abstract significantly more characters from the skull than the rest of  
547 the skeleton. However, tests for levels of homoplasy in the craniodental and postcranial  
548 partitions of our sample of 85 matrices revealed no significant differences, irrespective of  
549 how homoplasy was measured. Systematists appear to be coding characters of similar  
550 internal consistency from both regions of the body. It is unclear to what extent the bias  
551 towards coding craniodental characters reflects a real bias in their distribution, the  
552 availability of material, or arises from the preconceptions of systematists.
- 553 2. Craniodental and postcranial character partitions exhibited significant incongruence (ILD  
554 tests) in 31 of our 85 sample data sets. Likewise, our new  $IRD_{NND}$  test found significantly  
555 different relationships in 27 from 85 cases.
- 556 3. Although vertebrate systematists sometimes code morphological characters preferentially  
557 from particular anatomical regions (through necessity or by choice), received wisdom is that  
558 a broad and unbiased sampling is usually preferable when possible. Here, we provide



empirical evidence for this view. Our results strongly support dense sampling of characters from across the vertebrate skeleton.

4. The  $IRD_{NND}$  test appears to be less sensitive to differences in numbers of characters or amounts of missing data between partitions than the ILD test. We advocate its use as an ancillary test for addressing differences in implied relationships between data partitions.
5. There were significant differences in the distribution of significant partition inhomogeneity across higher taxonomic groups, as measured by the ILD test. Synapsids and amphibians were less likely to have congruent craniodental and postcranial partitions than other groups. However, these differences were no longer significant when the imbalance in partition size was modelled out. Neither were the relationships inferred from craniodental and postcranial characters more likely to conflict significantly in some higher taxa than in others ( $IRD_{NND}$  test). Our findings therefore have some generality across vertebrates.

## ACKNOWLEDGEMENTS

We would like to thank all authors who kindly supplied us with their datasets (knowingly or otherwise). We are grateful to Marcello Ruta and anonymous reviewers whose comments enabled us to substantially improve this paper. Ward Wheeler and Pablo Goloboff also made invaluable comments on an earlier version of this work as presented at *Hennig XXIX*. Some of our analyses were made possible by the generous free use of a high performance computing cluster; Bioportal, University of Oslo. RCPM gathered data and performed analyses. RS scripted  $IRD$  tests within *TNT* and performed analyses. MAW conceived and designed the study, devised/programmed the original tests in *PAUP\** and performed analyses. RCPM and MAW summarised results and wrote the paper. We thank Elizabeth Wills for drawing the silhouettes in Figure 3. This work was supported by a University of Bath URS award to RCPM, Leverhulme Trust Grant F/00351/Z and BBSRC grant BB/K015702/1 to MAW, JTF Grant 43915 to Mark Wilkinson and MAW, and NERC fellowship NE/I020253/1 to RSS.



589 **LITERATURE CITED**

- 590
- 591 Abourachid, A., E. Hofling. 2012. The legs: a key to bird evolutionary success. *J. Ornithol.* **153**: S193-S198.
- 592 Allain, R., and N. Aquesbi. 2008. Anatomy and phylogenetic relationships of *Tazoudasaurus naimi*
- 593 (Dinosauria, Sauropoda) from the late Early Jurassic of Morocco. *Geodiversitas* **30**: 345–424.
- 594 Allain, R., T. Xaisanavong, P. Richir, and B. Khentavong. 2012. The first definitive Asian spinosaurid
- 595 (Dinosauria: Theropoda) from the Early Cretaceous of Laos. *Naturwissenschaften* **99**: 369-377.
- 596 Allard, M. W., J. S. Farris, and J. M. Carpenter. 1999a. Congruence among mammalian mitochondrial genes.
- 597 *Cladistics* **15**: 75–84.
- 598 Allard, M. W., M. Kearney, K. L. Kivimaki, A. D. Meisner, E. E. Strong. 1999b. The random cladist: A review
- 599 of the software package *Random Cladistics*. *Cladistics* **15**: 183–189.
- 600 Alejandra Abello, M. 2013. Analysis of dental homologies and phylogeny of Paucituberculata (Mammalia:
- 601 Marsupialia). *Biol. J. Linn. Soc.* 109: 441-465.
- 602 Allen, V., K.T. Bates, Z.H. Li, J.R. Hutchinson. 2013. Linking the evolution of body shape and locomotor
- 603 biomechanics in bird-line archosaurs. *Nature* **497**: 104-107.
- 604 Anderson, J. S., R. R. Reisz, D. Scott, N. B. Frobisch, and S. S. Sumida. 2008. A stem batrachian from the
- 605 Early Permian of Texas and the origin of frogs and salamanders. *Nature* **453**: 515–518.
- 606 Andres, B., J. M. Clark, and X. Xing. 2010. A new rhamphorhynchid pterosaur from the Upper Jurassic of
- 607 Xinjiang China, and the phylogenetic relationships of basal pterosaurs. *J. Vertebr. Palaeontol.* **30**: 163-
- 608 187.
- 609 Archie, J. W. 1989. Homoplasy excess ratios: New indices for measuring levels of homoplasy in
- 610 phylogenetic systematics and a critique of the consistency index. *Syst. Biol.* **38**: 253–269.
- 611 Archie, J. W. 1996. Measures of homoplasy. Pp. 153-188 in M. J. Sanderson, L. Hufford (eds.) *Homoplasy:*
- 612 *The Recurrence of Similarity in Evolution*. Academic Press, New York.
- 613 Archie, J. W., and J. Felsenstein. 1993. The number of evolutionary steps on random and minimum length
- 614 trees for random evolutionary data. *Theor. Popul. Biol.* **43**: 52–79.
- 615 Arratia, G. 2009. Identifying patterns of diversity of the actinopterygian fulcra. *Acta Zoolog.* **90**: 220–235.
- 616 Asher, R. J. 2007. A web-database of mammalian morphology and a reanalysis of placental phylogeny. *BMC*
- 617 *Evol. Biol.* **7**: 108+.
- 618 Asher, R. J., and M. Hofreiter. 2006. Tenrec phylogeny and the noninvasive extraction of nuclear DNA. *Syst.*
- 619 *Biol.* **55**: 181–194.

620 Asher, R. J., and J. Müller. 2012. Molecular tools in palaeobiology: Divergence and mechanisms. Pp. 1-15 in  
621 R. J. Asher, and J. Müller (eds.) *From Clone to Bone: The Synergy of Morphological and Molecular*  
622 *Tools in Palaeobiology*. Cambridge University Press, Cambridge, UK.

623 Asher, R.J., J. Meng, J. R. Wible, M. C. McKenna, G. W. Rougier, D. Dashzeveg, M. J. Novacek. 2005. Stem  
624 Lagomorpha and the antiquity of Glires. *Science* **307**: 1091–1094.

625 Baker, R., X. Yu, and R. DeSalle. 1998. Assessing the relative contribution of molecular and morphological  
626 characters in simultaneous analysis trees. *Mol. Phylogenet. Evol.* **9**: 427–436.

627 Barker, F. K., and F. M. Lutzoni. 2002. The utility of the incongruence length difference test. *Syst. Biol.* **51**:  
628 625–37.

629 Baum, B. R. 1992. Combining trees as a way of combining data sets for phylogenetic inference, and the  
630 desirability of combining gene trees. *Taxon* **41**: 3–10.

631 Beard, K. C., L. Marivaux, Y. Chaimanee, J.-J. Jaeger, B. Marandat, P. Tafforeau, A. N. Soe, S. T. Tun, and  
632 A. A. Kyaw. 2009. A new primate from the Eocene Pondaung formation of Myanmar and the  
633 monophyly of Burmese amphipithecids. *Proc. Roy. Soc. Lond. B Biol. Sci.* **276**: 3285–3294.

634 Beck, R. M. D., H. Godthelp, V. Weisbecker, M. Archer, and S. J. Hand. 2008. Australia's oldest marsupial  
635 fossils and their biogeographical implications. *PLoS One* **3**: e1858+.

636 Billet, G. 2011. Phylogeny of the Notoungulata (Mammalia) based on cranial and dental characters. *J. Syst.*  
637 *Palaeontology* **9**: 481-497.

638 Bledsoe, A. H., and R. J. Raikow. 1990. A quantitative assessment of congruence between molecular and  
639 nonmolecular estimates of phylogeny. *J. Mol. Evol.* **30**: 247-259.

640 Bloch, J. I., M. T. Silcox, D. M. Boyer, and E. J. Sargis. 2007. New paleocene skeletons and the relationship  
641 of plesiadapiforms to crown-clade primates. *Proc. Natl. Acad. Sci. Unit. States Am.* **104**: 1159–1164.

642 Bluis, J., and D. G. Shin. 2003. Nodal distance algorithm: Calculating a phylogenetic tree comparison metric.  
643 Pp. 87-94 in *BIBE '03: Proceedings of the 3rd IEEE Symposium on Bioinformatics and*  
644 *BioEngineering*. IEEE Computer Society, Washington, DC.

645 Bouchard-Cote, A., S. Sankararaman, and M. I. Jordan. 2012. Phylogenetic inference via sequential Monte  
646 Carlo. *Syst. Biol.* **61**: 579-593.

647 Bourdon, E., A. Ricqlès, and J. Cubo. 2009. A new transantarctic relationship: Morphological evidence for a  
648 Rheidae–Dromaiidae–Casuariidae clade (Aves, Palaeognathae, Ratitae). *Zool. J. Linn. Soc.* **156**:  
649 641–662.

650 Bourque, M. 1978. Arbres de steiner et reseaux dont certains sommets sont a localisation variable. Thesis,  
651 University of Montreal.

652 Bremer, B. 1996. Combined and separate analyses of morphological and molecular data in the plant family  
653 Rubiaceae. *Cladistics* **12**: 21–40.

654 Brochu, C. A., J. Njau, R. J. Blumenschine, and L. D. Densmore. 2010. A new horned crocodile from the  
655 Plio-Pleistocene hominid sites at Olduvai Gorge, Tanzania. *PLoS One* **5**: e9333.

656 Bull, J. J., J. P. Huelsenbeck, C. W. Cunningham, D. L. Swofford, and P. J. Waddell. 1993. Partitioning and  
657 combining data in phylogenetic analysis. *Syst. Biol.* **42**: 384–397.

658 Burns, M. E., P. J. Currie, R. L. Sissons, and V. M. Arbour. 2011. Juvenile specimens of *Pinacosaurus*  
659 *grangeri* Gilmore, 1933 (Ornithischia: Ankylosauria) from the Late Cretaceous of China, with  
660 comments on the specific taxonomy of Pinacosaurus. *Cretaceous Research* **32**: 174-186.

661 Butler, R.J., P. Upchurch, and D. B. Norman. 2008. The phylogeny of the Ornithischian dinosaurs. *J. Syst.*  
662 *Palaeontology* **6**: 1–40.

663 Cardini, A., and S. Elton. 2008. Does the skull carry a phylogenetic signal? Evolution and modularity in the  
664 guenons. *Biol. J. Linn. Soc.* **93**: 813–834.

665 Cardini, A., and P.D. Polly. 2013. Larger mammals have longer faces because of size-related constraints on  
666 skull form. *Nature Comms.* **4**: 2458.

667 Carrano, M.T., and S. D. Sampson. 2008. The phylogeny of Ceratosauria (Dinosauria: Theropoda). *J. Syst.*  
668 *Palaeontology* **6**: 183–236.

669 Cau, A., and F. Fanti. 2011. The oldest known metriorhynchid crocodylian from the middle Jurassic of north-  
670 eastern Italy: *Neptunidraco ammoniticus* gen. et sp. nov. *Gondwana Research* **19**: 550-565.

671 Chase, K., D.R. Carrier, F.R. Adler, T. Jarvik, E.A. Ostrander, T.D. Lorentzen, and K.G. Lark. 2002. Genetic  
672 basis for systems of skeletal quantitative traits: Principal component analysis of the canid skeleton.  
673 *Proc. Natl. Acad. Sci. Unit. States. Am.* **99**: 9930-9935.

674 Cheng, L., X.-H. Chen, X.-W. Zeng, and Y.- J. Cai. 2012. A new eosauropterygian (Diapsida: Saurpterygia)  
675 from the Middle Triassic of Luoping, Yunnan Province. *J. Earth Sci.* **23**: 33-40.

676 Choo, B. 2011. Revision of the actinopterygian genus *Mimipiscis* (= *Mimia*) from the Upper Devonian Gogo  
677 Formation of Western Australia and the interrelationships of the early Actinopterygii. *Trans. Earth. Sci.*  
678 **102**: 77-104.

Clarke, J. A., and K. M. Middleton. 2008. Mosaicism, modules, and the evolution of birds: Results from a Bayesian approach to the study of morphological evolution using discrete character data. *Syst. Biol.* **57**: 185-201.

Cobbett A., M. Wilkinson, and M. A. Wills. 2007. Fossils impact as hard as living taxa in parsimony analyses of morphology. *Syst. Biol.* **56**: 753-766.

Collard, M., S. Gibbs, and B. Wood. 2001. Phylogenetic utility of higher primate postcranial morphology. *Am. J. Phys. Anthropol. Suppl.* **32**: 52.

Cunningham, C. W. 1997. Can three incongruence tests predict when data should be combined? *Mol. Biol. Evol.* **14**: 733–740.

de Pietri, V.L., L. Costeur, L. Guntert, and G. Mayr. 2011. A revision of the Lari (Aves, Charadriiformes) from the early Miocene of Saint-Gerand-le-Puy (Allier, France). *J. Vertebr. Paleontol.* **31**: 812-828.

de Vienne, D. M., T. Giraud, O. C. Martin. 2007. A congruence index for testing topological similarity between trees. *Bioinformatics (Oxford, England)* **23**: 3119–3124.

Degnan, J. H., and N. A. Rosenberg. 2009. Gene tree discordance, phylogenetic inference and the multispecies coalescent. *Trends Ecol. Evol.* **24**: 332–340.

Diogo, R. 2004. *Morphological Evolution, Adaptations, Homoplasies, Constraints, and Evolutionary Trends: Catfishes as a Case Study on General Phylogeny and Macroevolution*. Science Publishers, Enfield.

Dolphin, K., R. Belshaw, C. D. L. Orme, and D. L. J. Quicke. 2000. Noise and incongruence: Interpreting results of the incongruence length difference test. *Mol. Phylogenet. Evol.* **17**: 401–406.

Donoghue, M. J., and R. H. Ree. 2000. Homoplasy and developmental constraint: A model and an example from plants. *Am. Zool.* **40**: 759-769.

Draper, I., L. Hedenas, and G. L. Grimm. 2007. Molecular and morphological incongruence in European species of *Isoethecium* (Bryophyta). *Mol. Phylogenet. Evol.* **42**: 700–716.

Dutel, H., J. G. Maisey, D. R. Schwimmer, P. Janvier, M. Herbin, and G. Clement. 2012. The giant Cretaceous coelacanth (Actinistia, Sarcopterygii) *Megalocoelacanthus dobiei* Schwimmer, Stewart and Williams, 1994, and its bearing on Latimerioidei interrelationships. *PLoS One* **7**: DOI: 10.1371/journal.pone.0049911

Estabrook, G.F., F. R. McMorris, and C. A. Meacham. 1985. Comparison of undirected trees based on subtrees of four evolutionary units. *Syst. Zool.* **34**: 193–200.

708 Ezcurra, M. D., and G. Cuny. 2007. The coelophysoid *Lophostropheus airelensis*, gen. nov.: A review of the  
709 systematics of “*Liliensternus*” *airelensis* from the Triassic-Jurassic outcrops of Normandy (France). *J.*  
710 *Vertebr. Paleontol.* **27**: 73–86.

711 Faith, D. P., and P. S. Cranston. 1991. Could a cladogram this short have arisen by chance alone?: On  
712 permutation tests for cladistic structure. *Cladistics* **7**: 1–28.

713 Farke, A. A., M. J. Ryan, P. M. Barrett, D. H. Tanke, D. R. Braman, M. A. Loewen, and M. R. Graham. 2011.  
714 A new centrosaurine from the Late Cretaceous of Alberta, Canada, and the evolution of parietal  
715 ornamentation in horned dinosaurs. *Acta Palaeontologica Polonica* **56**: 691–702.

716 Farris, J. S. 1973. On comparing the shapes of taxonomic trees. *Syst. Zool.* **22**: 50–54.

717 Farris, J. S. 1989. The retention index and homoplasy excess. *Syst. Biol.* **38**: 406–407.

718 Farris, J. S., M. Källersjö, A. G. Kluge, and C. Bult. 1995a. Constructing a significance test for incongruence.  
719 *Syst. Biol.* **44**: 570–572.

720 Farris, J. S., M. Källersjö, A. G. Kluge, and C. Bult. 1995b. Testing significance of incongruence. *Cladistics*  
721 **10**: 315–319.

722 Felsenstein, J. 1988. Phylogenies from molecular sequences: Inference and reliability. *Annu. Rev. Genet.*  
723 **22**: 521–565.

724 Finarelli, J. A., and W. C. Clyde. 2004. Reassessing hominoid phylogeny: Evaluating congruence in the  
725 morphological and temporal data. *Paleobiology* **30**: 614–651.

726 Finden, C.R. and A.D. Gordon. 1985. Obtaining common pruned trees. *J. Classif.* **2**: 255–276.

727 Fitzgerald, E. M. G. 2010. The morphology and systematics of *Mammalodon colliveri* (Cetacea: Mysticeti), a  
728 toothed mysticete from the Oligocene of Australia. *Zool. J. Linn. Soc.* **158**: 367–476.

729 Frazzetta, T. H. 2012. Flatfishes, turtles, and bolyerine snakes: Evolution by small steps or large, or both?  
730 *Evol. Biol.* **39**: 30–60.

731 Freitas, A. V. L., and K. S. Brown. 2004. Phylogeny of the Nymphalidae (Lepidoptera). *Syst. Biol.* **53**: 363–  
732 383.

733 Friedman, M. 2007. *Styloichthys* as the oldest coelacanth: Implications for early osteichthyan  
734 interrelationships. *J. Syst. Palaeontology* **5**: 289–343.

735 Friedman, M. 2008. The evolutionary origin of flatfish asymmetry. *Nature* **454**: 209–212.

736 Frobisch, J. 2007. The cranial anatomy of *Kombuisia frerensis* Hotton (Synapsida, Dicynodontia) and a new  
737 phylogeny of anomodont therapsids. *Zool. J. Linn. Soc.* **150**: 117–144.

738 Gaffney, E. S., G. E. Hooks, and V.P. Schneider. 2009. New material of North American side-necked turtles  
739 (Pleurodira: Bothremydidae). *American Museum Novitates* **3655**: 1–26.

740 Gardiner, J.D., J.R. Codd, R.L. Nudds. 2011. An association between ear and tail morphologies of bats and  
741 their foraging style. *Can. J. Zool.* **89**: 90–99.

742 Gates, T.A., and S. D. Sampson. 2007. A new species of *Gryposaurus* (Dinosauria: Hadrosauridae) from the  
743 late Campanian Kaiparowits formation, Southern Utah, USA. *Zool. J. Linn. Soc.* **151**: 351–376.

744 Gatesy, J., and P. Arctander. 2000. Hidden morphological support for the phylogenetic placement of  
745 *Pseudoryx nghetinhensis* with bovine bovids: A combined analysis of gross anatomical evidence and  
746 DNA sequences from five genes. *Syst. Biol.* **49**: 515–538.

747 Gatesy, J., and R.H. Baker. 2005. Hidden likelihood support in genomic data: Can forty-five wrongs make a  
748 right? *Syst. Biol.* **54**: 483–492.

749 Gatesy, J., P. O’Grady, and R. H. Baker. 1999. Corroboration among data sets in simultaneous analysis:  
750 Hidden support for phylogenetic relationships among higher level artiodactyl taxa. *Cladistics* **15**: 271–  
751 313.

752 Gaubert, P., W. C. Wozencraft, P. Cordeiro-Estrela, and G. Veron. 2005. Mosaics of convergences and  
753 noise in morphological phylogenies: What’s in a viverrid-like carnivoran? *Syst. Biol.* **54**: 865–894.

754 Gaudin, T., R. Emry, and J. Wible. 2009. The phylogeny of living and extinct pangolins (Mammalia,  
755 Pholidota) and associated taxa: A morphology based analysis. *J. Mamm. Evol.* **16**: 235–305.

756 Goddard, W., E. Kubicka, G. Kubicki, and F. R. McMorris. 1994. The agreement metric for labelled binary  
757 trees. *Math. Biosci.* **123**: 215–226.

758 Godefroit, P., H. Shulin, Y. Tingxiang, and P. Lauters. 2008. New hadrosaurid dinosaurs from the uppermost  
759 Cretaceous of Northeastern China. *Acta Palaeontologica Polonica* **53**: 47–74.

760 Goloboff, P. A. 1999. Analyzing large data sets in reasonable times: Solutions for composite optima.  
761 *Cladistics* **15**: 415–428.

762 Goloboff, P. A. 2010. On weighting characters differently in different parts of the cladogram. *Cladistics* **26**:  
763 212.

764 Goloboff, P. A. 2008. Calculating SPR distances between trees. *Cladistics* **24**: 591–597.

765 Goloboff, P. A. 2014. Extended implied weighting. *Cladistics* **30**: 260–272.

766 Goloboff, P. A., J. M. Carpenter, J. S. Arias, and D. R. M. Esquivel. 2008a. Weighting against homoplasy  
767 improves phylogenetic analysis of morphological data sets. *Cladistics* **24**: 758–773.



768 Goloboff, P.A., J. S. Farris, and K. C. Nixon. 2008b. TNT, a free program for phylogenetic analysis. *Cladistics*  
769 **24**: 774-786.

770 Goswami, A., N. Milne, and S. Wroe. 2011. Biting through constraints: cranial morphology, disparity and  
771 convergence across living and fossil carnivorous mammals. *Proc. Roy. Soc. Lond. B Biol. Sci.* **278**:  
772 1831-1839.

773 Goswami, A., W.J. Binder, J. Meachen, and F.R. O'Keefe. 2015. The fossil record of phenotypic integration  
774 and modularity: A deep-time perspective on developmental and evolutionary dynamics. *Proc. Natl.*  
775 *Acad. Sci. Unit. States. Am.* **112**: 4891-4896.

776 Gould, G. C. 2001. The phylogenetic resolving power of discrete dental morphology among extant  
777 hedgehogs and the implications for their fossil record. *American Museum Novitates* **3340**: 1-52.

778 Gould, S. J. 1977. *Ontogeny and Phylogeny*. Belknap Press of Harvard University Press, Harvard.

779 Hamrick, M. W. 2012. The developmental origins of mosaic evolution in the primate limb skeleton. *Evol. Biol.*  
780 **39**: 447-455.

781 Hastings, A. K., J. I. Bloch, and C. A. Jaramillo. 2011. A new longirostrine dyrosaurid (Crocodylomorpha,  
782 Mesoeucrocodylia) from the Paleocene of north-eastern Colombia: Biogeographic and behavioural  
783 implications for New-World Dyrosauridae. *Palaeontology* **54**: 1095-1116.

784 Hill, R. V. 2005. Integration of morphological data sets for phylogenetic analysis of Amniota: The importance  
785 of integumentary characters and increased taxonomic sampling. *Syst. Biol.* **54**: 530–547.

786 Hillis, D. M., and J. J. Wiens. 2000. Molecules versus morphology in systematics: Conflicts, artifacts, and  
787 misconceptions. Pp. 1-19 in J. J. Wiens (ed.) *Phylogenetic Analysis of Morphological Data*.  
788 Smithsonian Institution Press, Washington DC.

789 Hilton, E. J., and P. L. Forey. 2009. Redescription of ?*Chondrosteus acipenseroides* Egerton, 1858  
790 (Acipenseriformes, ?Chondrosteidae) from the lower Lias of Lyme Regis (Dorset, England), with  
791 comments on the early evolution of sturgeons and paddlefishes. *J. Syst. Palaeontology* **7**: 427–453.

792 Hipp, A. L., J. C. Hall, and K. J. Sytsma, 2004. Congruence versus phylogenetic accuracy: Revisiting the  
793 incongruence length difference test. *Syst. Biol.* **53**: 81–89.

794 Holland, T., and J. A. Long. 2009. On the phylogenetic position of *Gogonasus andrewsae*, within the  
795 Tetrapodomorpha. *Acta Zoolog.* **90**: 285-296.

796 Hopkins, M. J, and S. Lidgard. 2012. Evolutionary mode routinely varies among morphological traits within  
797 fossil species lineages. *Proc. Natl. Acad. Sci. Unit. States. Am.* **50**: 20520-20525.

798 Hospitaleche, C. A., C. Tambussi, M. Donato, and M. Cozzuol. 2007. A new Miocene penguin from  
 799 Patagonia and its phylogenetic relationships. *Acta Palaeontologica Polonica* **52**: 299–314.  
 800 Hoyal-Cuthill, J.F., S. J. Braddy, and P. C. J. Donoghue. 2010. A formula for maximum possible steps in  
 801 multistate characters: Isolating matrix parameter effects on measures of evolutionary convergence.  
 802 *Cladistics* **26**: 98-102.  
 803 Hunt, G. 2008. Gradual or pulsed evolution: When should punctuational explanations be preferred?  
 804 *Paleobiology* **34**: 360-377.  
 805 Hurley, I. A., R. L. Mueller, K. A. Dunn, E. J. Schmidt, M. Friedman, R. K. Ho, V. E. Prince, Z. Yang, M. G.  
 806 Thomas, and M. I. Coates. 2007. A new time-scale for ray-finned fish evolution. *Proc. Roy. Soc. Lond.*  
 807 *B Biol. Sci.* **274**: 489–498.  
 808 Huson, D. H., and C Scornavacca. 2012. Dendroscope 3: An interactive tool for rooted phylogenetic trees  
 809 and networks. *Syst. Biol.* **61**: 1061-1067.  
 810 Imamura, H., S. M. Shirai, and M. Yabe. 2005. Phylogenetic position of the family Trichodontidae (Teleostei:  
 811 Perciformes), with a revised classification of the perciform suborder Cottoidei. *Ichthyological Research*  
 812 **52**: 264–274.  
 813 Jenner, R. A. 2004. When molecules and morphology clash: Reconciling conflicting phylogenies of the  
 814 Metazoa by considering secondary character loss. *Evol. Dev.* **6**: 372–378.  
 815 Ji, Q., Z.-X. Luo, and S.-A. Ji. 1999. A Chinese triconodon mammal and mosaic evolution of the mammalian  
 816 skeleton. *Nature* **398**: 326-330.  
 817 Ji, C., D.-Y. Jiang, R. Motani, W.-C. Hao, Z.-Y. Sun, and T. Cai. 2013. A new juvenile specimen of  
 818 *Guanlingsaurus* (Ichthyosauria, Shastasauridae) from the Upper Triassic of southwestern China. *J.*  
 819 *Vertebr. Paleontol.* **33**: 340-348.  
 820 Kangas, A. T., A. R. Evans, I. Thesleff, and J. Jernvall. 2004. Nonindependence of mammalian dental  
 821 characters. *Nature* **432**, 211-214.  
 822 Kemp, T. S. 2005. *The Origin and Evolution of Mammals*. Oxford University Press, Oxford.  
 823 Kemp, T. S. 2007. The concept of correlated progression as the basis of a model for the evolutionary origin  
 824 of major new taxa. *Proc. Roy. Soc. Lond. B Biol. Sci.* **274**: 1667-1673.  
 825 Kharlamova, A.V., L.N. Trut, D.R. Carrier, K. Chase and K.G. Lark. 2007. Genetic regulation of canine  
 826 skeletal traits: trade-offs between the hind limbs and forelimbs in the fox and dog. *Integr. Comp. Biol.*  
 827 **47**: 373–381.

828 Kitching, I., P. Forey, C. Humphries, and D. Williams. 1998. *Cladistics: Theory and Practice of Parsimony*  
829 *Analysis (Systematics Association Special Volume)*. Oxford University Press, Oxford.

830 Kivell, T. L., A. P. Barros, and J. B. Smaers. 2013. Different evolutionary pathways underlie the morphology  
831 of wrist bones in hominoids. *BMC Evol. Biol.* **13**: 1-12.

832 Kjaer, K. M., and R. L. Honeycutt. 2007. Site specific rates of mitochondrial genomes and the phylogeny of  
833 Eutheria. *BMC Evol. Biol.* **7**: 8.

834 Klassen, G. J., R. D. Mooi, and A. Locke. 1991. Consistency indices and random data. *Syst. Zool.* **40**: 446-  
835 457.

836 Klingenberg, C. P. 2008. Morphological integration and developmental modularity. *Annu. Rev. Ecol. Evol.*  
837 *Systemat.* **39**: 115-132.

838 Kluge, A. G. 1989. A concern for evidence and a phylogenetic hypothesis of relationships among *Epicrates*  
839 (Boidae, Serpentes). *Syst. Zool.* **38**: 7-25.

840 Kluge, A. G., and J. S. Farris. 1969. Quantitative phyletics and the evolution of anurans. *Syst. Zool.* **18**: 1-  
841 32.

842 Koski, K. 2007. The mandibular complex. *Eur. J. Orthod.* **29**: i118-i123.

843 Larouche, O., R. Cloutier, M.L Zelditch. 2015. Head, body and fins: patterns of morphological integration and  
844 modularity in fishes. *Evol. Biol.* **42**: 296-311.

845 Lee, M. S. Y., and A. B. Camens. 2009. Strong morphological support for the molecular evolutionary tree of  
846 placental mammals. *J. Evol. Biol.* **22**: 2243-2257.

847 Lee, M. S. Y., and J. D. Scanlon. 2002. Snake phylogeny based on osteology, soft anatomy and ecology.  
848 *Biol. Rev.* **77**: 333-401.

849 Lee, M. S. Y., and T. H. Worthy. 2012. Likelihood reinstates *Archaeopteryx* as a primitive bird. *Biol. Lett.* **8**:  
850 299-303.

851 Leighton, L. R., and C. G. Maples. 2002. Evaluating internal versus external characters: Phylogenetic  
852 analyses of the Echinoconchidae, Buxtoniinae, and Juresaniinae (Phylum Brachiopoda). *J. Paleontol.*  
853 **76**: 659-671.

854 Le Quesne, W. J. 1969. A method of selection of characters in numerical taxonomy. *Syst. Biol.* **18**: 201-205.

855 Lewis, P. 2000. A likelihood approach to estimating phylogeny from discrete morphological character data.  
856 *Syst. Biol.* **50**: 913-925.

857 Li, P.-P., K.-Q. Gao, L.-H. Hou, and X. Xu. 2007. A gliding lizard from the early Cretaceous of China. *Proc.*  
858 *Natl. Acad. Sci. Unit. States Am.* **104**: 5507-5509.

859 Lin, Y., V. Rajan, and B. Moret. 2011. A metric for phylogenetic trees based on matching bioinformatics  
860 research and applications. *Lect. Notes Comput. Sci.* **6674**: 197-208

861 Lister, A. M., C. J. Edwards, D. A. W. Nock, M. Bunce, I. A. van Pijlen, D. G. Bradley, M. G. Thomas, and I.  
862 Barnes. 2005. The phylogenetic position of the 'giant deer' *Megaloceros giganteus*. *Nature* **438**: 850–  
863 853.

864 Lloyd, G. T. 2009. Grame T. Lloyd Matrices. <http://www.graemetlloyd.com/matr.html>.

865 Lopez-Arbarello, A. 2011. Phylogenetic interrelationships of ginglymodian fishes (Actinopterygii:  
866 Neopterygii). *PLoS One* **7**: e39370. doi:10.1371/journal.pone.0039370.

867 Lü, J.-C., D. M. Unwin, X. Jin, Y. Liu, and Q. Ji. 2010. Evidence for modular evolution in a long-tailed  
868 pterosaur with a pterodactyloid skull. *Proc. Roy. Soc. Lond. B Biol. Sci.* **277**: 383–389.

869 Lü, J.-C., P. J. Currie, L. Xu, X.-L. Zhang, H.-Y. Pu, and S.-H. Jia. 2013. Chicken-sized oviraptorid dinosaurs  
870 from central China and their ontogenetic implications. *Naturwissenschaften* **100**: 165-175.

871 Lyson, T. R., and W. G. Joyce. 2009. A new species of *Palatobaena* (Testudines: Baenidae) and a maximum  
872 parsimony and Bayesian phylogenetic analysis of Baenidae. *J. Paleontol.* **83**, 457–470.

873 Maddison, W. P. 1997. Gene trees in species trees. *Syst. Biol.* **46**: 523–536.

874 Manegold, A., and T. Toepfer. 2013. The systematic position of *Hemicircus* and the stepwise evolution of  
875 adaptations for drilling, tapping and climbing up in true woodpeckers (Picinae, Picidae). *J. Zool.*  
876 *Systemat. Evol. Res.* **51**: 72-82.

877 Martinelli, A. G., and G. W. Rougier. 2007. On *Chaliminia musteloides* (Eucynodontia: Trithelodontidae) from  
878 the Late Triassic of Argentina, and a phylogeny of Ictidosauria. *J. Vertebr. Paleontol.* **27**: 442–460.

879 Martinez, R. N., and O. A. Alcober. 2009. A basal sauropodomorph (Dinosauria: Saurischia) from the  
880 Ischigualasto formation (Triassic, Carnian) and the early evolution of Sauropodomorpha. *PLoS One* **4**:  
881 e4397+.

882 Matsumoto, R., S. Suzuki, K. Tsogtbaatar, and S. Evans. 2009. New material of the enigmatic reptile  
883 *Khurendukhosaurus* (Diapsida: Choristodera) from Mongolia. *Naturwissenschaften* **96**: 233–242.

884 Mauricio, G. N., J. I. Areta, M. R. Bornschein, and R. E. Reis. 2012. Morphology-based phylogenetic  
885 analysis and classification of the family Rhinocryptidae (Aves: Passeriformes). *Zool. J. Linn. Soc.* **166**:  
886 377-432.

887 Maynard Smith, J. 1993. *The Theory of Evolution*. Cambridge University Press, Cambridge.

888 Mayr, G. 2009. Phylogenetic relationships of the paraphyletic caprimulgiform birds (nightjars and allies). *J.*  
889 *Zool. Syst. Evol. Res.* **48**: 126–137.

890 Mayr, G., R. S. Rana, K. D. Rose, A. Sahni, K. Kumar, L. Singh, and T. Smith. 2010. Quercypsitta-like birds  
891 from the early Eocene of India (Aves, ?Psittaciformes). *J. Vertebr. Paleontol.* **30**: 467-478.

892 Mayr, G. 2011. The phylogeny of charadriiform birds (shorebirds and allies) – reassessing the conflict  
893 between morphology and molecules. *Zool. J. Linn. Soc.* **161**: 916–934.

894 Mayr, G., and C. Mourer-Chauviré. 2004. Unusual tarsometatarsus of a mousebird from the Paleogene of  
895 France and the relationships of *Selmes* Peters, 1999. *J. Vertebr. Paleontol.* **24**: 366-372.

896 Mickevich, M. F. 1978. Taxonomic congruence. *Syst. Biol.* **27**: 143–158.

897 Mickevich, M. F., and J. S. Farris. 1981. The implications of congruence in *Menidia*. *Syst. Zool.* **30**: 351-370.

898 Mitteroecker, P., and F. Bookstein. 2007. The conceptual and statistical relationship between modularity and  
899 morphological integration. *Syst. Biol.* **56**: 818-836.

900 Mitteroecker, P., and F. Bookstein. 2008. The evolutionary role of modularity and integration in the hominoid  
901 cranium. *Evolution* **62**: 943-958.

902 Muller, J., and R. R. Reisz. 2006. The phylogeny of early eureptiles: Comparing parsimony and Bayesian  
903 approaches in the investigation of a basal fossil clade. *Syst. Biol.* **55**: 503–511.

904 Munoz-Duran, J. 2011. Data set incongruence, misleading characters, and insights from the fossil record:  
905 The canid phylogeny. *Caldasia* **33**: 637-658.

906 Murray, P.F., and P. Vickers-Rich. 2004. *Magnificent Mihirungs: The Colossal Flightless Birds of the*  
907 *Australian Dreamtime*. Indiana University Press, Bloomington.

908 Naylor, G. J. P., and D. C. Adams. 2001. Are the fossil data really at odds with the molecular data?  
909 Morphological evidence for Cetartiodactyla phylogeny reexamined. *Syst. Biol.* **50**: 444-453.

910 Nesbitt, S. J., N. D. Smith, R. B. Irmis, A. H. Turner, A. Downs, and M. A. Norell. 2009. A complete skeleton  
911 of a Late Triassic saurischian and the early evolution of dinosaurs. *Science* **326**: 1530-1533.

912 Nesbitt, S. J. 2011. The early evolution of archosaurs: Relationships and the origin of major clades. *Bull. Am.*  
913 *Mus. Nat. Hist.* **352**: 1-292.

914 Nesbitt, S. J., D. T. Ksepka, and J. A. Clarke . 2011. Podargiform affinities of the enigmatic *Fluvioviridavis*  
915 *platyrhamphus* and the early diversification of strisores (“Caprimulgiformes” + Apodiformes). *PLoS*  
916 *One* **6**: e26350.

917 Nichols, R. 2001. Gene trees and species trees are not the same. *Trends Ecol. Evol.* **16**: 358–364.

918 Nixon, K. C. 1999. The parsimony ratchet, a new method for rapid parsimony analysis. *Cladistics* **15**: 407-  
919 414.

920 Nylander, J. A. A, F. Ronquist, J. P. Huelsenbeck, and J. L. Nieves-Aldrey. 2003. Bayesian phylogenetic  
921 analysis of combined data. *Syst. Biol.* **53**: 47-67.

922 O'Connor, J.K., X. Wang, L.M. Chiappe, C. Gao, Q. Meng. 2009. Phylogenetic support for a specialized  
923 clade of Cretaceous Enantiornithine birds with information from a new species. *J. Vert. Paleontol.* **29**:  
924 188–204.

925 O'Connor, P. M., J. J. W. Sertich, N. J. Stevens, E. M. Roberts, M. D. Gottfried, T. L. Hieronymus, Z. A.  
926 Jinnah, R. Ridgely, S. E. Ngasala, and J. Temba. 2010. The evolution of mammal-like crocodyliforms  
927 in the Cretaceous period of Gondwana. *Nature* **466**: 748-751.

928 O'Leary, M.A., J. Gatesy, and M. J. Novacek. 2003. Are the dental data really at odds with the molecular  
929 data? Morphological evidence for whale phylogeny (Re)Reexamined. *Syst. Biol.* **52**: 853–864.

930 O'Leary, M. A., and S. Kaufman. 2011. MorphoBank: Phylophenomics in the "cloud". *Cladistics* **27**: 529-537.

931 O'Meara, B. 2009. Brian O'Meara Lab. <http://www.brianomeara.info/treebase>.

932 Osi, A., and L. Makadi. 2009. New remains of *Hungarosaurus tormai* (Ankylosauria, Dinosauria) from the  
933 Upper Cretaceous of Hungary: Skeletal reconstruction and body mass estimation. *Paleontologische*  
934 *Zeitschrift* **83**: 227–245.

935 Page, R. D. 1996. On consensus, confidence, and 'total evidence'. *Cladistics* **12**: 83–92.

936 Palci, A., M. W. Caldwell, and A. M. Albino. 2013. Emended diagnosis and phylogenetic relationships of the  
937 Upper Cretaceous fossil snake *Najash rionegrina* Apesteguía and Zaher, 2006. *Journal of Vertebrate*  
938 *Paleontology* **33**: 131-140.

939 Pamilo, P., M. Nei. 1988. Relationships between gene trees and species trees. *Mol. Biol. Evol.* **5**: 568–583.

940 Parenti, L. R. 2008. A phylogenetic analysis and taxonomic revision of ricefishes, *Oryzias* and relatives  
941 (Belontiiformes, Adrianichthyidae). *Zool. J. Linn. Soc.* **154**: 494–610.

942 Parrilla-Bel, J., M. T. Young, M. Moreno-Azanza, and J. I. Canudo. 2013. The first metriorhynchid  
943 crocodylomorph from the Middle Jurassic of Spain, with implications for evolution of the subclade  
944 Rhacheosaurini. *PLoS One* **8**: e54275+

945 Pattengale, N. D., E. J. Gottlieb, and B. M. E. Moret. 2007. Efficiently computing the Robinson-Foulds metric.  
946 *J. Comput. Biol.* **14**: 724–735.

947 Pattinson, D. J., R. S. Thompson, A. Piotrowski, and R. J. Asher. 2014. Phylogeny, palaeontology, and  
948 primates: Do incomplete fossils bias the tree of life? *Syst. Biol.* **64**: 169-86.

- 949 Penz, C. M., A. V. L. Freitas, L. A. Kaminski, M. M. Casagrande, and P. J. Devries. 2013. Adult and early-  
950 stage characters of Brassolini contain conflicting phylogenetic signal (Lepidoptera, Nymphalidae).  
951 *Syst. Entomol.* **38**: 316-333.
- 952 Phillips, M. J., T. H. Bennett, and M. S. Y. Lee. 2009. Molecules, morphology, and ecology indicate a recent,  
953 amphibious ancestry for echidnas. *Proc. Natl. Acad. Sci. Unit. States Am.* **106**: 17089–17094.
- 954 Pine, R. H., R. M. Timm, and M. Weksler. 2012. A newly recognized clade of trans-Andean oryzomyini  
955 (rodentia: Cricetidae), with description of a new genus. *J. Mammal.* **93**: 851-870.
- 956 Pisani, D., M. J. Benton, and M. Wilkinson. 2007. Congruence of morphological and molecular phylogenies.  
957 *Acta Biotheor.* **55**: 269-281.
- 958 Planet, P. 2006. Tree disagreement: Measuring and testing incongruence in phylogenies. *J. Biomed.*  
959 *Informat.* **39**: 86–102.
- 960 Planet, P. J., and I. N. Sarkarm. 2005. mILD: A tool for constructing and analyzing matrices of pairwise  
961 phylogenetic character incongruence tests. *Bioinformatics* **21**: 4423–4424.
- 962 Poe, S. 1996. Data set incongruence and the phylogeny of crocodilians. *Syst. Biol.* **45**: 393–414.
- 963 Pollitt, J. R., R. A. Fortey and M. A. Wills. 2015. Systematics of the trilobite families Lichidae Hawle & Corda,  
964 1847 and Lichakephalidae Tripp, 1957: The application of Bayesian inference to morphological data.  
965 *J. Syst. Palaeontology* **3**: 225-241.
- 966 Poyato-Ariza, F. J. 2003. Dental characters and phylogeny of pycnodontiform fishes. *J. Vertebr. Paleontol.*  
967 **23**: 937-940.
- 968 Puértolas, E., J. I. Canudo, and P. Cruzado-Caballero. 2011. A new crocodylian from the late Maastrichtian  
969 of Spain: Implications for the initial radiation of crocodyloids. *PLoS One* **6**: e20011+.
- 970 Pujos, F., G. de Iuliis, C. Argot, and L. Werdelin. 2007. A peculiar climbing Megalonychidae from the  
971 Pleistocene of Peru and its implication for sloth history. *Zool. J. Linn. Soc.* **149**: 179–235.
- 972 Quicke, D. L., O. R. Jones, and D. R. Epstein. 2007. Correcting the problem of false incongruence due to  
973 noise imbalance in the incongruence length difference (ILD) test. *Syst. Biol.* **56**: 496–503.
- 974 Rae, T. C. 1999. Mosaic evolution in the origin of the Hominoidea. *Folia Primatol.* **70**: 125-135.
- 975 Ragan, M. 1992. Matrix representation in reconstructing phylogenetic relationships among the eukaryotes.  
976 *Biosystems* **28**: 47–55.
- 977 Ramirez, M. J. 2006. Further problems with the incongruence length difference test: “Hypercongruence”  
978 effect and multiple comparisons. *Cladistics* **22**: 289–295.
- 979 Robinson, D. R., and L. R. Foulds. 1981. Comparison of phylogenetic trees. *Math. Biosci.* **53**: 131–147.

- Rodrigo A. G., M. Kelly-Borges, P. R. Bergquist, and P. L. Bergquist. 1993. A randomization test of the null hypothesis that two cladograms are sample estimates of a parametric phylogenetic tree. *New Zealand J. of Botany* **31**: 257-268.
- Rossello, F., and G. Valiente. 2006. The transposition distance for phylogenetic trees. *arXiv q-bio/0604024v1* [q-bio.PE]
- Ruta, M., and J. R. Bolt. 2008. The brachyopoid *Hadrokkosaurus bradyi* from the early Middle Triassic of Arizona, and a phylogenetic analysis of lower jaw characters in temnospondyl amphibians. *Acta Palaeontologica Polonica*, **53**: 579-592.
- Ruta, M., and M. I. Coates. 2007. Dates, nodes and character conflict: Addressing the lissamphibian origin problem. *J. Syst. Palaeontology* **5**: 69–122.
- Sánchez-Villagra, M. R., and B. A. Williams. 1998. Levels of homoplasy in the evolution of the mammalian skeleton. *J. Mamm. Evol.* **5**: 113–126.
- Sánchez-Villagra, M. R., I. Horovitz, and M. Motokawa. 2006. A comprehensive morphological analysis of talpid moles (Mammalia) phylogenetic relationships. *Cladistics* **22**: 59–88.
- Sanderson, M. J., and M. J. Donoghue. 1989. Patterns of variation in levels of homoplasy. *Evolution* **43**: 1781-1795.
- Sansom, R. S., S. E. Gabbott, and M. A. Purnell. 2010. Non-random decay of chordate characters causes bias in fossil interpretation. *Nature* **463**: 797-800.
- Sansom, R. S., S. E. Gabbott, and M. A. Purnell. 2011. Decay of vertebrate characters in hagfish and lamprey (Cyclostomata) and the implications for the vertebrate fossil record. *Proc. Roy. Soc. Lond. B Biol. Sci.* **1709**: 1150-1157.
- Sansom, R. S., and M. A. Wills. 2013. Fossilization causes organisms to appear erroneously primitive by distorting evolutionary trees. *Scientific Reports* **3**: 2545.
- Sansom, R. S. 2015. Bias and sensitivity in the placement of fossil taxa resulting from interpretations of missing data. *Syst. Biol.* **64**: 256-266.
- Sereno, P. C., and S. L. Brusatte. 2008. Basal abelisaurid and carcharodontosaurid theropods from the Lower Cretaceous Elrhaz formation of Niger. *Acta Palaeontologica Polonica* **53**: 15–46.
- Shimada, K. 2005. Phylogeny of lamniform sharks (Chondrichthyes: Elasmobranchii) and the contribution of dental characters to lamniform systematics. *Paleontol. Res.* **9**: 55–72.



- Sigurdson, T., A. K. Huttenlocker, S. P. Modesto, T. B. Rowe, and R. Damiani. 2012. Reassessment of the morphology and paleobiology of the therocephalian *Tetracynodon darti* (Therapsida), and the phylogenetic relationships of Baurioidea. *Journal of Vertebrate Paleontology* **32**: 1113-1134.
- Simmons, N. B., and T. M., Conway. 2001. Phylogenetic relationships of mormoopid bats (Chiroptera: Mormoopidae) based on morphological data. *Bull. Am. Mus. Nat. Hist.* **258**: 1–100.
- Simmons, N. B., K. L. Seymour, J. Habersetzer, and G. F. Gunnell. 2008. Primitive early Eocene bat from Wyoming and the evolution of flight and echolocation. *Nature* **451**: 818–821.
- Skutschas, P. P., and Y. M. Gubin. 2012. A new salamander from the late Paleocene—Early Eocene of Ukraine. *Acta Palaeontologica Polonica* **57**: 135-148.
- Smith, N. D., and D. Pol. 2007. Anatomy of a basal sauropodomorph dinosaur from the early Jurassic Hanson formation of Antarctica. *Acta Palaeontologica Polonica* **52**: 657–674.
- Smith, N. D. 2010. Phylogenetic analysis of Pelecaniformes (Aves) based on osteological data: Implications for waterbird phylogeny and fossil calibration studies. *PLoS One* **5**: e13354.
- Smith, N. 2011. Taxonomic revision and phylogenetic analysis of the flightless Mancallinae (Aves, Pan-Alcidae). *ZooKeys* **91**: 1-116.
- Sokal, R. R., and P. H. A. Sneath. 1963. *Principles of Numerical Taxonomy*. Freeman and Company, London.
- Song, H., and S. R. Bucheli. 2010. Comparison of phylogenetic signal between male genitalia and non-genital characters in insect systematics. *Cladistics* **26**: 23–35.
- Sparks, J. S. 2008. Phylogeny of the cichlid subfamily Etroplinae and taxonomic revision of the Malagasy cichlid genus *Paretroplus* (Teleostei: Cichlidae). *Bull. Am. Mus. Nat. Hist.* **314**: 1–151.
- Spaulding, M., M. A. O’Leary, and J. Gatesy. 2009. Relationships of Cetacea (Artiodactyla) among mammals: Increased taxon sampling alters interpretations of key fossils and character evolution. *PLoS One* **4**: e7062+.
- Springer, M.S., A. Burk-Herrick, R. Meredith, E. Eizirik, E. Teeling, S. J. O’Brien, and W. J. Murphy. 2007. The adequacy of morphology for reconstructing the early history of placental mammals. *Syst. Biol.* **56**: 673–684.
- Steel, M.A., and D. Penny. 1993. Distributions of tree comparison metrics – some new results. *Syst. Biol.* **42**: 126–141.

- Sues, H.-D., and A. Averianov. 2009. A new basal hadrosauroid dinosaur from the Late Cretaceous of Uzbekistan and the early radiation of duck-billed dinosaurs. *Proc. Roy. Soc. Lond. B Biol. Sci.* **276**: 2549–2555.
- Sues, H.-D., and R. R. Reisz. 2008. Anatomy and phylogenetic relationships of *Sclerosaurus armatus* (Amniota: Parareptilia) from the Buntsandstein (Triassic) of Europe. *J. Vertebr. Paleontol.* **28**: 1031–1042.
- Swartz, B. A. 2009. Devonian actinopterygian phylogeny and evolution based on a redescription of *Stegotrachelus finlayi*. *Zoo. J. Linn. Soc.* **156**: 750–784.
- Thompson, R. S., E. V. Barmann, and R. J. Asher. 2012. The interpretation of hidden support in combined data phylogenetics. *J. Zool. Syst. Evol. Res.* **50**: 251–263.
- Tsuji, L. A. 2013. Anatomy, cranial ontogeny and phylogenetic relationships of the pareiasaur *Deltavjatia rossicus* from the Late Permian of central Russia. *Earth and Environmental Science Transactions of the Royal Society of Edinburgh* **104**: 81–122.
- Tsuji, L. A., and J. Mueller. 2009. Assembling the history of the Parareptilia: phylogeny, diversification, and a new definition of the clade. *Fossil Record* **12**: 71–81.
- Turner, A. H., and J. J. W. Sertich. 2010. Phylogenetic history of *Simosuchus clarki* (Crocodyliformes: Notosuchia) from the late Cretaceous of Madagascar. *Journal of Vertebrate Paleontology* **30**: 177–236.
- Vallin, G., and M. Laurin. 2004. Cranial morphology and affinities of *Microbrachis*, and a reappraisal of the phylogeny and lifestyle of the first amphibians. *Journal of Vertebrate Paleontology* **24**: 56–72.
- Venczel, M. 2008. A new salamandrid amphibian from the Middle Miocene of Hungary and its phylogenetic relationships. *J. Syst. Palaeontology* **6**: 41–59.
- Vermeij, G. J., and S. J. Carlson. 2000. The muricid gastropod subfamily Rapaninae: Phylogeny and ecological history. *Paleobiology* **26**: 19–46.
- Vieira, G. H. C., G. R. Colli, and S. N. Bao. 2005. Phylogenetic relationships of corytophanid lizards (Iguania, Squamata, Reptilia) based on partitioned and total evidence analyses of sperm morphology, gross morphology, and DNA data. *Zoolog. Scripta* **34**: 605–625.
- Ward, C. V. 1997. Functional anatomy and phyletic implications of the hominoid trunk and hindlimb. Pp. 101–130 in B. Begun, C. Ward, and M. Rose (eds.). *Function, Phylogeny, and Fossils: Miocene Hominoid Evolution and Adaptations*. Plenum Press, New York.
- Waterman, M. S., and T. F. Smith. 1978. On the similarity of dendrograms. *J. Theor. Biol.* **73**: 789–800.

- Wheeler, W. 1999. Measuring topological congruence by extending character techniques. *Cladistics* **15**: 131–135.
- Wiens, J. J. 2003a. Incomplete taxa, incomplete characters, and phylogenetic accuracy: Is there a missing data problem? *J. Vertebr. Paleontol.* **23**: 297–310.
- Wiens, J. J. 2003b. Missing data, incomplete taxa, and phylogenetic accuracy. *Syst. Biol.* **52**: 528–538.
- Wiens, J. J., and B. D. Hollingsworth. 2000. War of the iguanas: Conflicting molecular and morphological phylogenies and long-branch attraction in iguanid lizards. *Syst. Biol.* **49**: 143–159.
- Wilkinson, M. 1995. TAXEQ3: Safe taxonomic reduction using taxonomic equivalence. Computer program published by the author. <http://www.softpedia.com/get/Science-CAD/TAXEQ3.shtml>
- Williams, W. T., and H. T. Clifford. 1971. On the comparison of two classifications of the same set of elements. *Taxon* **20**: 519–522.
- Wills, M. A., R. A. Jenner, and C. Ni Dhubhghaill. 2009. Eumalacostracan evolution: conflict between three sources of data. *Arthropod Systematics and Phylogeny* **67**: 71–90.
- Wilson, M. V. H., and A. M. Murray. 2008. Osteoglossomorpha: phylogeny, biogeography, and fossil record and the significance of key African and Chinese fossil taxa. Pp. 185–219 in L. Cavin, A. Longbottom, and M. Richter (eds.). *Fishes and the Break-up of Pangaea*. Geological Society of London, Special Publications.
- Xu, G.-H., and K.-Q. Gao, 2011. A new scanilepiform from the Lower Triassic of northern Gansu Province, China, and phylogenetic relationships of non-teleostean Actinopterygii. *Zoo. J. Linn. Soc.* **161**: 595–612.
- Young, M. T., and M. B. de Andrade. 2009. What is *Geosaurus*?: Redescription of *Geosaurus giganteus* (Thalattosuchia: Metriorhynchidae) from the Upper Jurassic of Bayern, Germany. *Zool. J. Linn. Soc.* **157**: 551–585.
- Young, N. M. 2005. Estimating hominoid phylogeny from morphological data: Character choice, phylogenetic signal and postcranial data. Pp. 19–31 in D. E. Lieberman, R. J. Smith, and J. Kelley (eds.) *Interpreting the Past: Essays on Human, Primate and Mammal Evolution in Honor of David Pilbeam*. Brill Academic Publishers Inc., Boston.
- Zanno, L. E., D. D. Gillette, L. B. Albright, and A. L. Titus. 2009. A new North American therizinosaurid and the role of herbivory in ‘predatory’ dinosaur evolution. *Proc. Roy. Soc. Lond. B Biol. Sci.* **276**: 3505–3511.

1098     Zaretskii, K. A. 1965. Constructing a tree on the basis of a set of distances between the hanging vertices.  
1099             *Uspekhi Mat. Nauk* **20**: 90–92.  
1100  
1101  
1102  
1103



| Author(s)               | Year | Clade                | Taxa | Cran. chars. | Post. chars. | Cran. miss % | Post. miss % | ILD   | IRD NND | Cran. CI | Post. CI | Cran. ci | Post. ci | Cran. HER | Post. HER |
|-------------------------|------|----------------------|------|--------------|--------------|--------------|--------------|-------|---------|----------|----------|----------|----------|-----------|-----------|
| Bourdon et al.          | 2009 | Palaeognathae        | 17   | 35           | 94           | 4.0          | 8.0          | 1.000 | 0.979   | 0.956    | 0.855    | 0.971    | 0.908    | 0.975     | 0.921     |
| Clarke & Middleton      | 2008 | Avialae              | 20   | 45           | 141          | 38.7         | 22.7         | 0.001 | 0.127   | 0.740    | 0.643    | 0.774    | 0.732    | 0.734     | 0.703     |
| de Pietri et al.        | 2011 | Charadriiformes      | 12   | 15           | 25           | 0.6          | 2.3          | 0.158 | 0.685   | 0.667    | 0.758    | 0.567    | 0.756    | 0.383     | 0.442     |
| Hospitaleche et al.     | 2007 | Sphenisciformes      | 19   | 20           | 23           | 5.5          | 14.6         | 0.290 | 0.600   | 0.479    | 0.509    | 0.539    | 0.617    | 0.502     | 0.367     |
| Manegold & Toepfer      | 2013 | Picidae              | 27   | 20           | 47           | 1.7          | 0.4          | 0.485 | 0.005   | 0.710    | 0.458    | 0.816    | 0.647    | 0.904     | 0.761     |
| Mauricio et al.         | 2012 | Rhynocriptidae       | 38   | 40           | 46           | 1.1          | 1.1          | 0.043 | 0.048   | 0.480    | 0.517    | 0.646    | 0.654    | 0.721     | 0.710     |
| Mayr et al.             | 2010 | Basal Psittaciformes | 32   | 25           | 68           | 3.9          | 3.3          | 0.466 | 0.180   | 0.456    | 0.341    | 0.541    | 0.456    | 0.400     | 0.355     |
| Mayr                    | 2010 | Caprimulgiformes     | 10   | 18           | 51           | 1.7          | 2.4          | 0.572 | 0.374   | 0.692    | 0.643    | 0.735    | 0.735    | 0.604     | 0.522     |
| Mayr                    | 2011 | Pelagornithidae      | 25   | 19           | 39           | 21.3         | 8.1          | 0.002 | 0.071   | 0.431    | 0.388    | 0.569    | 0.419    | 0.494     | 0.426     |
| Nesbitt et al.          | 2011 | Strisores            | 20   | 36           | 61           | 13.6         | 23.1         | 0.002 | 0.002   | 0.656    | 0.447    | 0.756    | 0.566    | 0.753     | 0.379     |
| O'Connor et al.         | 2009 | Enantiornithines     | 19   | 40           | 152          | 43.7         | 29.7         | 0.067 | 0.557   | 0.712    | 0.578    | 0.760    | 0.699    | 0.608     | 0.565     |
| Smith                   | 2010 | Pelecaniformes       | 53   | 94           | 370          | 5.3          | 2.7          | 0.845 | 0.865   | 0.493    | 0.441    | 0.622    | 0.581    | 0.804     | 0.799     |
| Smith                   | 2011 | Mancallinae          | 54   | 51           | 163          | 3.5          | 6.7          | 0.063 | 0.175   | 0.214    | 0.212    | 0.313    | 0.331    | 0.511     | 0.581     |
| Allain & Aquesbi        | 2008 | Sauropods            | 18   | 62           | 150          | 31.1         | 21.5         | 0.006 | 0.009   | 0.683    | 0.686    | 0.756    | 0.790    | 0.662     | 0.682     |
| Allain et al.           | 2012 | Spinosaurids         | 23   | 75           | 94           | 19.5         | 24.7         | 0.044 | 0.025   | 0.514    | 0.579    | 0.649    | 0.675    | 0.516     | 0.656     |
| Andres et al.           | 2010 | Basal Pterosaurs     | 18   | 44           | 31           | 18.2         | 17.7         | 0.054 | 0.556   | 0.667    | 0.574    | 0.718    | 0.649    | 0.658     | 0.537     |
| Burns et al.            | 2011 | Ankylosaurs          | 17   | 45           | 15           | 12.8         | 36.9         | 0.365 | 0.024   | 0.548    | 0.704    | 0.661    | 0.661    | 0.527     | 0.439     |
| Butler et al.           | 2008 | Ornithischia         | 29   | 124          | 88           | 19.8         | 25.0         | 0.367 | 0.134   | 0.502    | 0.521    | 0.666    | 0.693    | 0.574     | 0.604     |
| Carrano & Sampson       | 2008 | Ceratosauria         | 12   | 66           | 76           | 36.3         | 31.9         | 0.304 | 0.451   | 0.863    | 0.769    | 0.924    | 0.846    | 0.854     | 0.690     |
| Ezcurra & Cuny          | 2007 | Coelophysoidea       | 11   | 63           | 73           | 12.8         | 15.3         | 0.011 | 0.791   | 0.610    | 0.731    | 0.678    | 0.768    | 0.617     | 0.668     |
| Gates & Sampson         | 2007 | Hadrosaurinae        | 13   | 59           | 14           | 14.5         | 20.3         | 0.276 | 0.024   | 0.607    | 0.667    | 0.679    | 0.674    | 0.499     | 0.390     |
| Godefroit et al.        | 2008 | Hadrosauridae        | 15   | 30           | 7            | 9.3          | 8.5          | 1.000 | 0.003   | 0.918    | 1.000    | 0.958    | 0.956    | 0.927     | 1.000     |
| Lu et al.               | 2010 | Pterosauria          | 37   | 59           | 55           | 8.5          | 21.0         | 0.002 | 0.066   | 0.561    | 0.457    | 0.695    | 0.572    | 0.698     | 0.624     |
| Lu et al.               | 2013 | Oviraptoridae        | 17   | 108          | 69           | 31.1         | 33.2         | 0.002 | 0.001   | 0.756    | 0.516    | 0.837    | 0.581    | 0.760     | 0.315     |
| Martinez & Alcober      | 2009 | Dinosauria           | 12   | 28           | 70           | 33.6         | 27.7         | 0.067 | 0.779   | 0.732    | 0.560    | 0.780    | 0.677    | 0.544     | 0.251     |
| Nesbitt et al.          | 2009 | Early Dinosaurs      | 35   | 113          | 194          | 32.2         | 30.8         | 0.001 | 0.003   | 0.413    | 0.469    | 0.540    | 0.603    | 0.429     | 0.653     |
| Nesbitt                 | 2011 | Archosaurs           | 55   | 166          | 223          | 23.1         | 28.2         | 0.001 | 0.030   | 0.402    | 0.413    | 0.576    | 0.562    | 0.592     | 0.706     |
| Osi & Makadi            | 2009 | Ankylosauria         | 15   | 43           | 14           | 20.3         | 32.9         | 0.117 | 0.216   | 0.537    | 0.643    | 0.626    | 0.598    | 0.502     | 0.827     |
| Sereno                  | 2008 | Carcharodontosaurids | 9    | 37           | 23           | 31.8         | 29.5         | 0.485 | 0.327   | 0.848    | 0.639    | 0.899    | 0.692    | 0.950     | 0.201     |
| Smith & Pol             | 2007 | Plateosauria         | 26   | 115          | 232          | 23.2         | 17.9         | 0.250 | 0.414   | 0.461    | 0.362    | 0.550    | 0.533    | 0.563     | 0.581     |
| Sues & Averianov        | 2009 | Iguanodontidae       | 25   | 100          | 38           | 11.9         | 24.4         | 0.265 | 0.014   | 0.650    | 0.672    | 0.280    | 0.305    | 0.838     | 0.827     |
| Zanno et al.            | 2009 | Therizinosauria      | 21   | 80           | 191          | 29.3         | 25.1         | 0.054 | 0.380   | 0.528    | 0.524    | 0.617    | 0.609    | 0.460     | 0.540     |
| Asher et al.            | 2005 | Lagomorpha           | 29   | 157          | 58           | 7.1          | 1.8          | 0.001 | 0.048   | 0.373    | 0.331    | 0.460    | 0.403    | 0.501     | 0.429     |
| Asher & Hofreiter       | 2006 | Afrotheria           | 20   | 90           | 22           | 9.0          | 4.4          | 0.795 | 0.617   | 0.406    | 0.441    | 0.487    | 0.445    | 0.404     | 0.325     |
| Asher                   | 2007 | Eutheria             | 46   | 121          | 75           | 6.4          | 5.2          | 0.001 | 0.040   | 0.223    | 0.275    | 0.305    | 0.406    | 0.286     | 0.384     |
| Beard et al.            | 2009 | Amphiphihcedidae     | 22   | 216          | 55           | 17.0         | 37.6         | 0.004 | 0.478   | 0.305    | 0.569    | 0.503    | 0.555    | 0.426     | 0.427     |
| Beck et al.             | 2008 | Marsupiala           | 27   | 101          | 139          | 10.7         | 9.1          | 0.004 | 0.259   | 0.432    | 0.400    | 0.532    | 0.517    | 0.543     | 0.477     |
| Bloch et al.            | 2007 | Plesiadapiforms      | 12   | 98           | 61           | 3.2          | 17.3         | 0.001 | 0.062   | 0.401    | 0.627    | 0.531    | 0.702    | 0.337     | 0.295     |
| Gaubert et al.          | 2005 | Feliformia           | 39   | 229          | 100          | 1.1          | 3.1          | 0.001 | 0.500   | 0.356    | 0.335    | 0.266    | 0.219    | 0.497     | 0.519     |
| Gaudin et al.           | 2009 | Pholidota            | 16   | 298          | 90           | 14.8         | 23.3         | 0.528 | 0.155   | 0.339    | 0.325    | 0.598    | 0.658    | 0.522     | 0.532     |
| Lister et al.           | 2005 | Cervidae             | 10   | 21           | 41           | 0.5          | 1.7          | 0.729 | 0.410   | 0.750    | 0.623    | 0.778    | 0.659    | 0.418     | 0.261     |
| Martinelli & Rougier    | 2007 | Ichthyosauria        | 16   | 65           | 17           | 10.9         | 31.3         | 0.088 | 0.001   | 0.674    | 0.770    | 0.790    | 0.778    | 0.646     | 0.837     |
| Phillips et al.         | 2009 | Monotremes           | 55   | 325          | 126          | 25.3         | 21.0         | 0.004 | 0.002   | 0.337    | 0.503    | 0.543    | 0.625    | 0.682     | 0.811     |
| Pine et al.             | 2012 | Oryzomyini           | 36   | 55           | 34           | 3.1          | 16.5         | 0.033 | 0.207   | 0.271    | 0.384    | 0.371    | 0.388    | 0.248     | 0.304     |
| Pujos et al.            | 2007 | Foliwora             | 18   | 25           | 17           | 7.8          | 21.6         | 0.066 | 0.441   | 0.455    | 0.628    | 0.503    | 0.610    | 0.382     | 0.457     |
| Sanchez-Villagra et al. | 2006 | Talpidae             | 17   | 73           | 83           | 4.7          | 4.3          | 0.003 | 0.010   | 0.495    | 0.549    | 0.587    | 0.652    | 0.445     | 0.609     |
| Sigurdson et al.        | 2012 | Baurioidea           | 20   | 95           | 20           | 8.3          | 35.8         | 0.507 | 0.049   | 0.520    | 0.706    | 0.638    | 0.725    | 0.588     | 0.597     |
| Simmons & Conway        | 2011 | Mormoopidae          | 16   | 103          | 106          | 16.0         | 14.0         | 1.000 | 0.974   | 0.650    | 0.659    | 0.539    | 0.586    | 0.679     | 0.685     |
| Simmons et al.          | 2008 | Chiroptera           | 25   | 48           | 154          | 16.9         | 21.7         | 0.201 | 0.559   | 0.445    | 0.419    | 0.524    | 0.521    | 0.525     | 0.459     |
| Spaulding et al.        | 2009 | Artiodactyla         | 37   | 353          | 137          | 17.6         | 21.0         | 0.018 | 0.711   | 0.235    | 0.307    | 0.326    | 0.444    | 0.288     | 0.444     |
| Brochu et al.           | 2010 | Crocodyloidea        | 24   | 57           | 23           | 7.3          | 14.1         | 0.032 | 0.030   | 0.598    | 0.540    | 0.703    | 0.530    | 0.658     | 0.464     |
| Cheng et al.            | 2012 | Sauropterygia        | 35   | 69           | 70           | 10.7         | 14.0         | 0.463 | 0.830   | 0.401    | 0.405    | 0.503    | 0.471    | 0.471     | 0.586     |
| Gaffney et al.          | 2009 | Bothremydidae        | 23   | 100          | 50           | 18.0         | 27.1         | 0.001 | 0.186   | 0.686    | 0.727    | 0.789    | 0.800    | 0.733     | 0.729     |
| Hill                    | 2005 | Amniota              | 19   | 124          | 150          | 1.4          | 4.6          | 0.457 | 0.616   | 0.550    | 0.620    | 0.655    | 0.729    | 0.664     | 0.737     |
| Ji et al.               | 2013 | Ichthyosauria        | 23   | 42           | 56           | 27.4         | 16.9         | 0.024 | 0.001   | 0.620    | 0.716    | 0.655    | 0.808    | 0.653     | 0.822     |
| Lee & Scanlon           | 2002 | Serpentes            | 21   | 204          | 53           | 9.9          | 4.3          | 0.882 | 0.209   | 0.505    | 0.522    | 0.627    | 0.569    | 0.548     | 0.472     |
| Li et al.               | 2007 | Squamata             | 23   | 212          | 97           | 8.1          | 15.1         | 0.099 | 0.389   | 0.536    | 0.511    | 0.644    | 0.614    | 0.591     | 0.447     |
| Lyson & Joyce           | 2009 | Baenidae             | 10   | 31           | 17           | 11.2         | 14.7         | 0.960 | 0.155   | 0.672    | 0.708    | 0.753    | 0.755    | 0.503     | 0.571     |
| Matsumoto et al.        | 2009 | Choristodera         | 14   | 49           | 29           | 14.6         | 13.3         | 0.002 | 0.728   | 0.647    | 0.571    | 0.704    | 0.558    | 0.611     | 0.424     |
| Muller & Reisz          | 2006 | Eureptiles           | 24   | 58           | 19           | 12.3         | 16.3         | 0.171 | 0.028   | 0.418    | 0.468    | 0.479    | 0.309    | 0.347     | 0.399     |
| Palci et al.            | 2013 | Serpentes            | 26   | 181          | 29           | 15.4         | 13.4         | 0.613 | 0.945   | 0.468    | 0.595    | 0.594    | 0.639    | 0.570     | 0.662     |
| Parilla-Bel et al.      | 2013 | Metrirhynchoidea     | 29   | 139          | 59           | 25.0         | 38.3         | 0.191 | 0.005   | 0.582    | 0.862    | 0.692    | 0.883    | 0.743     | 0.918     |
| Sues & Reisz            | 2008 | Parareptilia         | 12   | 34           | 9            | 10.3         | 38.8         | 0.034 | 0.004   | 0.731    | 0.750    | 0.797    | 0.657    | 0.738     | 0.631     |
| Tsuji & Mueller         | 2013 | Pareiasauria         | 21   | 64           | 50           | 22.3         | 36.3         | 0.118 | 0.066   | 0.755    | 0.793    | 0.811    | 0.847    | 0.772     | 0.795     |
| Vieira et al.           | 2005 | Corytophanidae       | 12   | 40           | 28           | 0.8          | 0.3          | 0.485 | 0.625   | 0.788    | 0.673    | 0.829    | 0.745    | 0.819     | 0.597     |
| Anderson et al.         | 2008 | Batrachia            | 52   | 143          | 76           | 18.4         | 29.3         | 0.001 | 0.911   | 0.254    | 0.285    | 0.347    | 0.367    | 0.407     | 0.379     |
| Holland & Long          | 2009 | Tetrapodomorpha      | 12   | 54           | 41           | 8.5          | 21.3         | 0.009 | 0.093   | 0.800    | 0.710    | 0.843    | 0.774    | 0.777     | 0.655     |
| Ruta & Coates           | 2007 | Lissamphibia         | 46   | 189          | 104          | 20.9         | 15.7         | 0.001 | 0.196   | 0.320    | 0.314    | 0.466    | 0.470    | 0.460     | 0.463     |
| Skutchas & Gubin        | 2012 | Salamanders          | 21   | 16           | 40           | 18.5         | 34.2         | 0.067 | 0.015   | 0.565    | 0.491    | 0.512    | 0.558    | 0.283     | 0.276     |
| Vallin & Laurin         | 2004 | Stegocephalia        | 47   | 99           | 60           | 15.8         | 26.7         | 0.002 | 0.028   | 0.355    | 0.417    | 0.435    | 0.538    | 0.584     | 0.544     |
| Venczel                 | 2008 | Caudata              | 15   | 19           | 16           | 1.8          | 12.5         | 1.000 | 0.145   | 0.538    | 0.783    | 0.596    | 0.854    | 0.516     | 0.876     |
| Choo                    | 2011 | Actinopterygii       | 27   | 51           | 26           | 12.4         | 16.4         | 0.635 | 0.460   | 0.424    | 0.448    | 0.550    | 0.600    | 0.464     | 0.451     |
| Dutel et al.            | 2012 | Actinistia           | 30   | 83           | 23           | 29.1         | 22.1         | 0.904 | 0.288   | 0.424    | 0.500    | 0.566    | 0.548    | 0.523     | 0.442     |
| Friedman                | 2007 | Actinistia           | 25   | 136          | 40           | 25.0         | 36.6         | 0.003 | 0.150   | 0.464    | 0.523    | 0.631    | 0.691    | 0.580     | 0.511     |
| Friedman                | 2008 | Pleuronectiformes    | 19   | 16           | 42           | 19.4         | 8.9          | 0.284 | 0.223   | 0.548    | 0.526    | 0.599    | 0.595    | 0.423     | 0.608     |
| Hilton & Forey          | 2009 | Acipenseriformes     | 18   | 31           | 17           | 6.5          | 6.5          | 0.661 | 0.010   | 0.634    | 0.724    | 0.751    | 0.740    | 0.670     | 0.739     |
| Hurley et al.           | 2007 | Actinopterygii       | 24   | 53           | 16           | 19.3         | 17.4         | 0.484 | 0.001   | 0.506    | 0.562    | 0.632    | 0.606    | 0.562     | 0.421     |
| Inamura et al.          | 2005 | Cottoidei            | 8    | 16           | 15           | 0.0          | 0.0          | 0.353 | 0.273   | 0.640    | 0.700    | 0.656    | 0.810    | 0.390     | 0.601     |
| Lopez-Arbarello         | 2011 | Ginglymodi           | 29   | 65           | 19           | 24.0         | 15.4         | 0.321 | 0.071   | 0.454    | 0.471    | 0.612    | 0.489    | 0.576     | 0.477     |
| Parenti                 | 2008 | Adrianichthyidae     | 31   | 37           | 43           | 0.6          | 8.8          | 0.574 | 0.565   | 0.677    | 0.584    | 0.844    | 0.644    | 0.778     | 0.688     |
| Shimada                 | 2005 | Lamniiformes         | 16   | 40           | 15           | 12.7         | 1.3          | 0.214 | 0.284   | 0.525    | 0.810    | 0.552    | 0.769    | 0.551     | 0.860     |

1106  
1107  
1108  
1109  
1110  
1111  
1112  
1113  
1114  
1115  
1116  
1117  
1118  
1119  
1120  
1121  
1122

**Table 1.** Summary statistics for the 85 vertebrate morphological data sets analysed herein. Data set dimensions (numbers of taxa (ntax) and informative characters (nchar)) refer to the pre-processed matrices after applying safe taxonomic deletion rules (see text for details). ‘Cran. char.’ and ‘Post. char’ denote characters per partition. ‘Cran. miss %’ and ‘Post. miss %’ report the percentage of missing data cells for partitions. ‘ILD’ column reports the p-value resulting from an incongruence length difference test with 999 random partitions. IRD columns report the p-values resulting from incongruence relationship difference tests with 999 random partitions. ‘IRD<sub>NND</sub>’ denotes the results of the IRD test using the Robinson Foulds tree-to-tree distance for nearest neighbouring trees. CI columns give ensemble consistency indices for partitions of the data set (craniodental or postcranial). Similarly, ci columns report the mean per character consistency indices for partitions of the data set (craniodental or postcranial) when optimised onto the globally most parsimonious trees. HER gives the homoplasy excess ratio for partitions of the data set (craniodental or postcranial) derived from 999 randomized matrices. An expanded version of this table containing additional statistics is provided within the Supporting Information.

1123

| Taxonomic Group                | Aves | Other<br>Ornithodira | Synapsida | Other<br>Reptiles | Amphibians | Fishes | Log<br>likelihood<br>ratio (G) | <i>p</i> |
|--------------------------------|------|----------------------|-----------|-------------------|------------|--------|--------------------------------|----------|
| No. data sets                  | 13   | 19                   | 18        | 15                | 6          | 14     |                                |          |
| Consistency indices sig. diff. | 8    | 8                    | 9         | 7                 | 4          | 4      | 4.283                          | 0.51     |
| ILD                            | 4    | 7                    | 10        | 5                 | 4          | 1      | 11.808                         | 0.038    |
| IRD NND                        | 3    | 9                    | 6         | 5                 | 2          | 2      | 4.801                          | 0.441    |
| IRD MR                         | 1    | 7                    | 5         | 3                 | 0          | 1      | 9.522                          | 0.09     |

1124


1125

1126

1127

**Table 2.** Number of data matrices in higher taxonomic groups, with a tally of those with significant ( $p < 0.05$ ) results for various partition homogeneity tests. G-test and  $p$  values quoted for the null that significant results are equally likely in the five higher taxonomic groups in each case.






**Figure 1.** (a and b) Characters sampled from different anatomical regions can yield radically different most parsimonious trees (MPT) when analysed in isolation. In both cases, there is no homoplasy *within* either region (characters 1-3 or characters 4-6), and a single MPT results in each case. (c) Combining the data from both partitions (characters 1-6) yields four MPTs, the strict consensus of which (illustrated) is entirely unresolved. Character statistics have been averaged over the four trees. (d) Two additional characters (5' and 6') are sampled from the same region as 'b', and these have the same distribution as 5 and 6 respectively. Analysis of all characters now reveals a single MPT with relationships identical to those in 'b' (characters 4-6). Characters 4-6, 5' and 6' contain no homoplasy: all conflicts are resolved with a cost to characters 1-3. In this case, the MPT is identical to the result that would be obtained by a clique analysis (*sensu* Le Quesne 1969).

## Craniodental partition


## Postcranial partition



**Figure 2.** Calculation of two partition inhomogeneity metrics for 'craniodental' and 'postcranial' partitions of a hypothetical data set. In this example there are equal numbers of craniodental (1-9) and postcranial (10-18) characters, but this need not be the case. For the Incongruence Length Difference (ILD) measure, maximally parsimonious trees (MPTs) are inferred from the craniodental and postcranial partitions of the data *independently*. The summed lengths of these trees (11 steps + 12 steps) is the sum of partition lengths (23 steps). In parallel with this, an MPT is inferred from both partitions analysed *simultaneously*. This tree is longer (25 steps) than the sum of partition lengths (23 steps), and the difference between them is the ILD (25 - 23 = 2). The ILD represents the reduction in homoplasy afforded by the isolation of the two partitions (two extra steps are needed when the partitions are combined). For the Incongruence Relationship Difference (IRD) measure, the branching structure of the craniodental and postcranial partition MPTs are compared (rather than their lengths) using one of several possible tree-to-tree distance metrics. Here, we illustrate the symmetric difference distance (RF) of Robinson and Foulds (1991). Open circles mark branches in either the craniodental or postcranial MPT that are absent from the other. The tally of these unique branches on both trees is the RF (2 + 2 = 4). Some background level of ILD or IRD is anticipated wherever a data set contains homoplasy. In order to interpret these observed metrics, therefore, we need to know what values would be expected for partitions of similar data sets in similar proportions. Random character partitions are used to

1158 generate null distributions for both the ILD and IRD, and observed values deemed significantly different from  
 1159 the null if they lie in some specified fraction of the tails.

1160




1161

1162

1163 **Figure 3.** Most parsimonious trees derived from craniodental and postcranial partitions can be significantly  
 1164 more different than we would expect. Tanglegrams computed using *Dendroscope* (Huson and Scornavacca,

2012). (a) The theropod data of Ezcurra and Cuny (2007) yielded one most parsimonious tree from the craniodental partition and six from the postcranial partition, the latter summarized as a majority rule tree merely for ease of visualization (we advocate the use of tests based upon mean nearest neighbours within sets of most parsimonious trees). Branches labelled with circles are unique to one or other tree (those unlabelled are common to both). The Robinson Foulds (RF) distance between the two is simply the sum of unique branches (6+6=12). While the ILD test returned a highly significant result ( $p = 0.011$ ) and our new incongruence relationship difference test (IRD<sub>NND</sub>) using RF did not ( $p = 0.791$ ). (b) Six craniodental and four postcranial trees in the mammalian data of Beck (2008), again summarized as majority rule consensus trees for visualization. In this case, the incongruence length difference (ILD) test for partition homogeneity returned a highly significant result ( $p = 0.004$ ) whereas our incongruence relationship difference (IRD<sub>NND</sub>) test did not ( $p = 0.259$ ). Indicative images are, from top to bottom on right hand side: *Ornithorhynchus*, *Tachyglossus*, *Didelphis*, *Monodelphis*, *Caenolestes*, *Dasyurus*, *Phascolarctos*, *Vombatus*, *Perameles*, *Echymipera*, *Macropus*, *Phalanger*, *Petaurus*, *Vincelestes*. Image of *Tachyglossus* courtesy of echidnasclub.com. See text for further explanation.


**Robinson Foulds Distances**



**Figure 4.** Summarising sets of most parsimonious trees (MPTs) for partitions prior to calculating tree-to-tree distances is computationally much faster than calculating distances between nearest neighbours. However, majority rule trees present modal or most frequent relationships, and may therefore plot eccentricly in tree

space. This is an undesirable property when attempting to summarise distances between sets of trees.

Figure shows tree-to-tree distances for craniodental and postcranial partitions of the mammalian data of Pujos (2007). Distance matrices have been plotted in two dimensions using non-metric multidimensional scaling (NMDS), and rotated using principal components analysis (PCA). Circles indicate craniodental trees and squares indicate postcranial trees. Open symbols denote original MPTs, filled symbols (black) denote majority rule trees.



**Figure 5.** Summary of results from three partition homogeneity tests, subdivided by higher taxonomic group. Black bars denote significant ( $p < 0.05$ ) differences in ci values between partitions (Mann-Whitney U tests). Gray bars denote significant ILD results, while light gray bars indicate significant IRD<sub>NND</sub> results. Images from phylopic.org (courtesy of Michael Scroggie, Oscar Sanisidro, B. Kimmel and Steven Coombs). Salamander and pterosaur courtesy of Matt Reinbold (modified by T. Michael Keesey) and Mark Witton. (<http://creativecommons.org/licenses/by-sa/3.0/>).